

## Datasheet: MCA5953F

<b>Description:</b>	MOUSE ANTI BOVINE CD21:FITC
<b>Specificity:</b>	CD21
<b>Other names:</b>	CR2
<b>Format:</b>	FITC
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	CC51
<b>Isotype:</b>	IgG2b
<b>Quantity:</b>	0.1 mg

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10
Immunofluorescence			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

#### Target Species

Bovine

#### Species Cross Reactivity

Reacts with: Pig, African Buffalo

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

#### Product Form

Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

#### Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
FITC	490	525

#### Preparation

Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> ) 1% Bovine Serum Albumin
<b>Approx. Protein Concentrations</b>	IgG concentration 0.1 mg/ml
<b>Immunogen</b>	Bovine (Friesian cattle) mesenteric lymph node cells.
<b>External Database Links</b>	<b>UniProt:</b> <a href="#">Q8HY44</a> <a href="#">Related reagents</a>
<b>Fusion Partners</b>	Spleen cells from immunized BALB/c mice were fused with cells of the mouse NSI myeloma cell line
<b>Specificity</b>	<p><b>Mouse anti Bovine CD21 monoclonal antibody, clone CC51</b> recognizes the bovine homologue of the human CD21 cell surface antigen, a 145 kDa single pass type I membrane glycoprotein containing multiple <a href="#">sushi</a> domains. CD21 is also known as complement receptor type 2 (CR2). In cattle CD21 expression is restricted to B-cells (<a href="#">Naessens et al. 1990</a>). CD21 may be expressed on B-cells as either a long or a short form (<a href="#">Pringle et al. 2012</a>).</p> <p>Mouse anti Bovine CD21, clone CC51 demonstrates cross reactivity with porcine and provides a reliable marker for porcine B-Cells (<a href="#">Sinkora et. al. 2013</a>). In addition to clone CC51, clone CC21 (<a href="#">MCA1424GA</a>) which has been demonstrated to recognise CD21 in a range of ruminant and other species is also available from Bio-Rad.</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul
<b>References</b>	<ol style="list-style-type: none"> <li>Denham S. <i>et al.</i> (1994) Monoclonal antibodies recognising differentiation antigens on porcine B cells. <a href="#">Vet Immunol Immunopathol. 43: 259-67.</a></li> <li>Boersma W.J. <i>et al.</i> (2001) Summary of workshop findings for porcine B-cell markers. <a href="#">Vet Immunol Immunopathol. 80: 63-78.</a></li> <li>Tenaya I.W. <i>et al.</i> (2012) Flow cytometric analysis of lymphocyte subset kinetics in Bali cattle experimentally infected with Jembrana disease virus. <a href="#">Vet Immunol Immunopathol. 149: 167-76.</a></li> <li>Sinkora, M <i>et al.</i> (2013) Different anti-CD21 antibodies can be used to discriminate developmentally and functionally different subsets of B lymphocytes in circulation of pigs. <a href="#">Dev Comp Immunol. 39: 409-18.</a></li> <li>Sinkora, M. <i>et al.</i> (2014) The comparative profile of lymphoid cells and the T and B cell spectratype of germ-free piglets infected with viruses SIV, PRRSV or PCV2. <a href="#">Vet Res. 45: 91.</a></li> <li>Liu, J. <i>et al.</i> (2020) <i>Theileria annulata</i>. transformation altered cell surface molecules expression and endocytic function of monocyte-derived dendritic cells. <a href="#">Ticks Tick Borne Dis. 11 (3): 101365.</a></li> <li>Li, J. <i>et al.</i> (2023) Single-cell transcriptomic analysis reveals transcriptional and cell</li> </ol>

subpopulation differences between human and pig immune cells. [Genes Genomics. Nov 18 \[Epub ahead of print\].](#)

8. Seemann, L. *et al.* (2024) Dietary L-carnitine supplementation modifies blood parameters of mid-lactating dairy cows during standardized lipopolysaccharide-induced inflammation. [Front Immunol. 15: 1390137.](#)

9. Yuan, C. *et al.* (2024) Effects of porcine epidemic diarrhea virus infection on CD21(+) B cells activation. [Vet Microbiol. 293: 110087.](#)

---

**Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA5953F>  
10041

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG2b NEGATIVE CONTROL:FITC \(MCA691F\)](#)

### Recommended Useful Reagents

[MOUSE ANTI PIG CD3:RPE \(MCA5951PE\)](#)

[MOUSE ANTI PIG CD3 \(MCA5951GA\)](#)

[MOUSE ANTI PIG SLA CLASS II DR \(MCA2314GA\)](#)

[RAT ANTI HUMAN CD3:Pacific Blue® \(MCA1477PB\)](#)

[MOUSE ANTI BOVINE CD4:RPE \(MCA1653PE\)](#)

[MOUSE ANTI HUMAN CD14:Pacific Blue® \(MCA1568PB\)](#)

[MOUSE ANTI HUMAN CD14:RPE \(MCA1568PE\)](#)

[MOUSE ANTI BOVINE CD25:RPE \(MCA2430PE\)](#)

[MOUSE ANTI BOVINE MHC CLASS II DR:RPE \(MCA5656PE\)](#)

[MOUSE ANTI PIG CD4 ALPHA:RPE \(MCA1749PE\)](#)

[MOUSE ANTI PIG CD25 \(MCA1736GA\)](#)

[MOUSE ANTI PIG wCD8 ALPHA:RPE \(MCA1223PE\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M385305:210513'

