

Datasheet: MCA5954F

Description:	MOUSE ANTI PIG CD8 BETA CHAIN:FITC
Specificity:	CD8 BETA
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	PPT23
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/10
Immunofluorescence			■	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Pig		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide (NaN ₃)		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml		
Immunogen	Porcine thymus membrane lysate		
Fusion Partners	Lymph node cells from immunized BALB/c mice were fused with cells of the NSO myeloma cell line		
Specificity	Mouse anti Pig CD8 beta chain, clone PPT23 , recognizes the porcine homologue of the human CD8 beta chain cell surface antigen.		

Characterization of clone PPT23, also known under the clone designation FYP1C5, has demonstrated that on peripheral blood lymphocytes and spleen, this antibody binds to cells that are CD3⁺, CD4⁻, CD8^{hi} and as such defines this antibody as a specific marker of porcine α/β T cells. Characterization of clone PPT23 has shown that in thymic tissue both CD8^{lo} and CD8^{hi} cells are recognized (Yang H & Parkhouse R.M. 1997). Inhibition studies have demonstrated that clone PPT23 recognizes a different epitope of on the CD8 beta chain to clone PPT22.

Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul
References	<ol style="list-style-type: none">1. Yang, H. & Parkhouse, R.M. (1997) Differential expression of CD8 epitopes amongst porcine CD8-positive functional lymphocyte subsets. Immunology. 92 (1): 45-52.2. Mair, K.H. <i>et al.</i> (2015) Carbopol improves the early cellular immune responses induced by the modified-live vaccine Ingelvac PRRS[®] MLV. Vet Microbiol. 176 (3-4): 352-7.3. Sinkora, M. <i>et al.</i> (2007) Two groups of porcine TCRgammadelta+ thymocytes behave and diverge differently. J Immunol. 178 (2): 711-9.
Further Reading	<ol style="list-style-type: none">1. Zuckermann, F.A. <i>et al.</i> (1998) Report on the analyses of mAb reactive with porcine CD8 for the second international swine CD workshop. Vet Immunol Immunopathol. 60 (3-4): 291-303.
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	18 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

Recommended Useful Reagents

[MOUSE ANTI PIG CD3:RPE \(MCA5951PE\)](#)

[MOUSE ANTI PIG CD4 ALPHA:RPE \(MCA1749PE\)](#)

[MOUSE ANTI PIG CD4 ALPHA \(MCA1749GA\)](#)

[MOUSE ANTI PIG CD3 \(MCA5951GA\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M334768:181203'

Printed on 11 Oct 2019