

## Datasheet: MCA5960GA

**BATCH NUMBER 151476**

|                      |  |
|----------------------|--|
| <b>Description:</b>  | MOUSE ANTI HUMAN SPHINGOSINE 1- PHOSPHATE RECEPTOR 1 |
| <b>Specificity:</b>  | SPHINGOSINE 1- PHOSPHATE RECEPTOR 1                  |
| <b>Other names:</b>  | CD363, S1P1  |
| <b>Format:</b>       | Purified   |
| <b>Product Type:</b> | Monoclonal Antibody                                  |
| <b>Clone:</b>        | 2B9  |
| <b>Isotype:</b>      | IgG2a  |
| <b>Quantity:</b>     | 0.1 mg   |

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

|                            | Yes | No | Not Determined | Suggested Dilution |
|----------------------------|-----|----|----------------|--------------------|
| Flow Cytometry             | ▪   |    |                | 1/10               |
| Immunohistology - Frozen   |     |    | ▪              |                    |
| Immunohistology - Paraffin |     |    | ▪              |                    |
| ELISA                      |     |    | ▪              |                    |
| Immunoprecipitation        | ▪   |    |                | 1/300              |
| Western Blotting           | ▪   |    |                | 1/5000             |
| Immunofluorescence         | ▪   |    |                | 1/500              |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

|                                 |  |
|---------------------------------|--|
| <b>Target Species</b>           | Human  |
| <b>Product Form</b>             | Purified IgG - liquid  |
| <b>Preparation</b>              | Purified IgG prepared by affinity chromatography on Protein A from ascites |
| <b>Buffer Solution</b>          | Phosphate buffered saline  |
| <b>Preservative Stabilisers</b> | 0.09% Sodium Azide (NaN <sub>3</sub> )                                     |

|                                       |   |
|---------------------------------------|---|
| <b>Approx. Protein Concentrations</b> | IgG concentration 1.0 mg/ml   |
| <b>External Database Links</b>        | <p><b>UniProt:</b><br/> <a href="#">P21453</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b><br/> <a href="#">1901</a>    S1PR1    <a href="#">Related reagents</a></p>  |
| <b>Synonyms</b>                       | CHEDG1, EDG1  |
| <b>Specificity</b>                    | <p><b>Mouse anti Human Sphingosine 1-Phosphate Receptor 1, clone 2B9</b> recognizes S1P1, otherwise known as CD363/EDG-1, a bioactive lysophospholipid and inducible member of the G-protein coupled receptor superfamily (GPCR) which plays an important role in a variety of cellular processes.</p> <p>During inflammatory diseases such as atherosclerosis the phenotype switching of activated macrophages is influenced by the presence of particular cytokines within the microenvironment. These cells have been characterized as classically activated M1 macrophages, in response to pro-inflammatory molecules such as interferon gamma and lipopolysaccharide, or as alternatively activated M2 macrophages, in response to anti-inflammatory cytokines such as interleukin-4. Activation of S1P1 through its ligand S1P (Sphingosine 1-Phosphate) has been identified as an important promoter of anti-inflammatory M2 macrophages (<a href="#">Hughes et al. 2008</a>).</p> <p>Apoptotic cells can also influence the pro-versus anti-inflammatory macrophage response, and S1P acts as a positive regulator for anti-apoptotic/pro-survival responses by binding to S1P1. Growing evidence suggests that the expression of S1P1 on macrophages may convey protection against apoptosis, at least in part, having implications for the pathogenesis of inflammatory diseases and cancer.</p> <p>Deviation in sphingosine 1-phosphate (S1P) signaling is important in many types of cancer. Analysis of S1P1 and SK1 receptors during the development of tamoxifen resistance in Estrogen Receptor (ER) breast cancer patients concurs that the expression levels of S1P1 correlates with tumor size and Progesterone Receptor (PR) status and that high S1P1 membrane expression is associated with shorter time to recurrence.</p> |
| <b>Flow Cytometry</b>                 | Use 10ul of the suggested working dilution to label $1 \times 10^6$ cells in 100ul  |
| <b>Storage</b>                        | <p>Store at +4°C or at -20°C if preferred.</p> <p>Storage in frost-free freezers is not recommended.</p> <p>This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>  |
| <b>Guarantee</b>                      | 12 months from date of despatch   |

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at:  
<https://www.bio-rad-antibodies.com/SDS/MCA5960GA>  
10040

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)  
Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)  
Goat Anti Mouse IgG (STAR76...) [RPE](#)  
Goat Anti Mouse IgG (STAR70...) [FITC](#)  
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)  
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)  
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)  
Goat Anti Mouse IgG (STAR77...) [HRP](#)  
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),  
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),  
[FITC](#), [HRP](#)

### Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

|                                  |   |                  |   |               |   |
|----------------------------------|---|------------------|---|---------------|---|
| <b>North &amp; South America</b> | Tel: +1 800 265 7376<br>Fax: +1 919 878 3751<br>Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a> | <b>Worldwide</b> | Tel: +44 (0)1865 852 700<br>Fax: +44 (0)1865 852 739<br>Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a> | <b>Europe</b> | Tel: +49 (0) 89 8090 95 21<br>Fax: +49 (0) 89 8090 95 50<br>Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a> |
|----------------------------------|---|------------------|---|---------------|---|

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M368527:200529'

Printed on 17 Mar 2025

---

© 2025 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)