

## Datasheet: MCA596APCT

<b>Description:</b>	MOUSE ANTI HUMAN CD14:APC
<b>Specificity:</b>	CD14
<b>Format:</b>	APC
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	UCHM1
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	25 TESTS/0.25ml

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

### Target Species

Human

### Species Cross Reactivity

Reacts with: Cynomolgus monkey, Rhesus Monkey, Fish, Trout  
**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

### Product Form

Purified IgG conjugated to Allophycocyanin (APC) - lyophilized

### Reconstitution

Reconstitute with 0.25 ml distilled water

### Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
APC	650	661

### Preparation

Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide 1% Bovine Serum Albumin 5% Sucrose
<b>Immunogen</b>	Human Thymocytes followed by peripheral blood mononuclear cells.
<b>External Database Links</b>	<b>UniProt:</b> <a href="#">P08571</a> <a href="#">Related reagents</a>  <b>Entrez Gene:</b> <a href="#">929</a> CD14 <a href="#">Related reagents</a>
<b>RRID</b>	AB_1100705
<b>Fusion Partners</b>	Spleen cells from immunized BALB/c mice were fused with cells from the NS1-Ag4/1 mouse myeloma line.
<b>Specificity</b>	<b>Mouse anti Human CD14 antibody, clone UCHM1</b> recognizes a cell surface antigen of ~55 kDa, known as CD14. The CD14 molecule is found predominantly on monocytes and macrophages in flow cytometry, it is less strongly expressed on granulocytes, and is absent from stem cells and myeloid cells of very early differentiation states. In immunohistology the CD14 molecule is found to be present on Langerhans cells, follicular dendritic cells, histiocytes and high endothelial venules. Antibodies to the CD14 molecule are known to induce oxidative burst formation. In tonsil tissue sections UCHM1 gives positive staining reactions with monocytic cells, the interfollicular tissue macrophages seen under the capsule, and dendritic reticulum cells. Skin Langerhans cells are always negative ( <a href="#">Hogg <i>et al.</i> 1984</a> ). UCHM1 also reacts with Kupffer cells and sinus lining cells on the liver.
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>Hogg, N. <i>et al.</i> (1984) Monoclonal antibodies specific for human monocytes, granulocytes and endothelium. <a href="#">Immunology. 53 (4): 753-67.</a></li> <li>Linch, D.C. <i>et al.</i> (1984) Monoclonal antibodies differentiating between monocytic and nonmonocytic variants of AML. <a href="#">Blood. 63 (3): 566-73.</a></li> <li>Angel, C.E. <i>et al.</i> (2006) Cutting edge: CD1a+ antigen-presenting cells in human dermis respond rapidly to CCR7 ligands. <a href="#">J Immunol. 176 (10): 5730-4.</a></li> <li>Köller, M. <i>et al.</i> (2004) Phenotypic and functional deficiencies of monocyte-derived dendritic cells in systemic lupus erythematosus (SLE) patients. <a href="#">Int Immunol. 16: 1595-604.</a></li> <li>Kämmerer, U. <i>et al.</i> (2003) Unique appearance of proliferating antigen-presenting cells expressing DC-SIGN (CD209) in the decidua of early human pregnancy. <a href="#">Am J Pathol. 162: 887-96.</a></li> <li>Goddard, S. <i>et al.</i> (2004) Interleukin-10 secretion differentiates dendritic cells from human liver and skin. <a href="#">Am J Pathol. 164: 511-9.</a></li> <li>Bournazos, S. <i>et al.</i> (2008) Monocyte functional responsiveness after PSGL-1-mediated</li> </ol>

- platelet adhesion is dependent on platelet activation status. [Arterioscler Thromb Vasc Biol. 28: 1491-8.](#)
8. Hovden, A.O. *et al.* (2011) Maturation of monocyte derived dendritic cells with OK432 boosts IL-12p70 secretion and conveys strong T-cell responses. [BMC Immunol. 12:2.](#)
  9. Hsu, T.L. *et al.* (2002) Modulation of dendritic cell differentiation and maturation by decoy receptor 3. [J Immunol. 168: 4846-53.](#)
  10. Karlsson, H. *et al.* (2002) Innate immune responses of human neonatal cells to bacteria from the normal gastrointestinal flora. [Infect Immun. 70: 6688-96.](#)
  11. Din JN *et al.* (2013) Effect of  $\omega$ -3 fatty acid supplementation on endothelial function, endogenous fibrinolysis and platelet activation in male cigarette smokers. [Heart. 99 \(3\): 168-74.](#)
  12. Fischer, U. and Koellner, B. (2007) Cross-reactivity of human leukocyte differentiation antigen monoclonal antibodies on carp and rainbow trout cells. [Vet Immunol Immunopathol. 119: 142-55.](#)
  13. Lin, C.W. *et al.* (2005) CD94 1A transcripts characterize lymphoblastic lymphoma/leukemia of immature natural killer cell origin with distinct clinical features. [Blood. 106: 3567-74.](#)
  14. Angel, C.E. *et al.* (2009) Distinctive localization of antigen-presenting cells in human lymph nodes. [Blood. 113: 1257-67.](#)
  15. Chang, Y.C. *et al.* (2004) Modulation of macrophage differentiation and activation by decoy receptor 3. [J Leukoc Biol. 75: 486-94.](#)
  16. Iking-Konert, C. *et al.* (2008) T lymphocytes in patients with primary vasculitis: expansion of CD8+ T cells with the propensity to activate polymorphonuclear neutrophils. [Rheumatology \(Oxford\). 47: 609-16.](#)
  17. Brook, F.A. *et al.* (2010) Derivation and characterisation of the human embryonic stem cell line, OxF1. [In Vitro Cell Dev Biol Anim. 46: 173-7.](#)
  18. Bromberek, J.L. *et al.* (2016) Breed Distribution and Clinical Characteristics of B Cell Chronic Lymphocytic Leukemia in Dogs. [J Vet Intern Med. 30 \(1\): 215-22.](#)
  19. Amouroux, R. *et al.* (2016) *De novo* DNA methylation drives 5hmC accumulation in mouse zygotes. [Nat Cell Biol. 18 \(2\): 225-33.](#)
  20. Spiller, K.L. *et al.* (2016) Differential gene expression in human, murine, and cell line-derived macrophages upon polarization. [Exp Cell Res. 347 \(1\): 1-13.](#)
  21. Kannegieter, N.M. *et al.* (2018) Analysis of NFATc1 amplification in T cells for pharmacodynamic monitoring of tacrolimus in kidney transplant recipients. [PLoS One. 13 \(7\): e0201113.](#)
  22. Matsusaka, K. *et al.* (2022) Distinct roles in phagocytosis of the early and late increases of cell surface calreticulin induced by oxaliplatin [Biochem Biophys Res. 29: 101222.](#)
  23. Wu, T.C. *et al.* (2018) IL1 Receptor Antagonist Controls Transcriptional Signature of Inflammation in Patients with Metastatic Breast Cancer. [Cancer Res. 78 \(18\): 5243-58.](#)
  24. Hoang, P.T. *et al.* (2018) Subtype Diversification and Synaptic Specificity of Stem Cell-Derived Spinal Interneurons. [Neuron. 100 \(1\): 135-149.e7.](#)

---

**Storage**

Prior to reconstitution store at +4°C.

After reconstitution store at +4°C.

DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we

recommend microcentrifugation before use.

---

<b>Guarantee</b>	12 months from date of despatch
------------------	---------------------------------

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #20487 available at: 20487: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/20487.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/20487.pdf</a>
--------------------------------------	--

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL:APC \(MCA929APC\)](#)

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M391213:211008'

Printed on 06 May 2022