

Datasheet: MCA596GT

BATCH NUMBER 150392

Description:	MOUSE ANTI HUMAN CD14
Specificity:	CD14
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	UCHM1
Isotype:	IgG2a
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50
Immunohistology - Frozen (1)	▪			
Immunohistology - Paraffin		▪		
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

Target Species	Human
Species Cross Reactivity	Reacts with: Cynomolgus monkey, Rhesus Monkey, Fish, Trout N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Human Thymocytes followed by peripheral blood mononuclear cells.
External Database Links	<p>UniProt: P08571 Related reagents</p> <p>Entrez Gene: 929 CD14 Related reagents</p>
RRID	AB_1100701
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells from the NS1-Ag4/1 mouse myeloma line.
Specificity	<p>Mouse anti Human CD14 antibody, clone UCHM1 recognizes a cell surface antigen of ~55 kDa, known as CD14. The CD14 molecule is found predominantly on monocytes and macrophages in flow cytometry, it is less strongly expressed on granulocytes, and is absent from stem cells and myeloid cells of very early differentiation states. In immunohistology the CD14 molecule is found to be present on Langerhans cells, follicular dendritic cells, histiocytes and high endothelial venules. Antibodies to the CD14 molecule are known to induce oxidative burst formation. In tonsil tissue sections UCHM1 gives positive staining reactions with monocytic cells, the interfollicular tissue macrophages seen under the capsule, and dendritic reticulum cells. Skin Langerhans cells are always negative (Hogg et al. 1984). UCHM1 also reacts with Kupffer cells and sinus lining cells on the liver.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
Histology Positive Control Tissue	Tonsil
References	<ol style="list-style-type: none"> Hogg, N. <i>et al.</i> (1984) Monoclonal antibodies specific for human monocytes, granulocytes and endothelium. Immunology. 53 (4): 753-67. Linch, D.C. <i>et al.</i> (1984) Monoclonal antibodies differentiating between monocytic and

- nonmonocytic variants of AML. [Blood. 63 \(3\): 566-73.](#)
3. Angel, C.E. *et al.* (2006) Cutting edge: CD1a+ antigen-presenting cells in human dermis respond rapidly to CCR7 ligands. [J Immunol. 176 \(10\): 5730-4.](#)
 4. Köller, M. *et al.* (2004) Phenotypic and functional deficiencies of monocyte-derived dendritic cells in systemic lupus erythematosus (SLE) patients. [Int Immunol. 16: 1595-604.](#)
 5. Kämmerer, U. *et al.* (2003) Unique appearance of proliferating antigen-presenting cells expressing DC-SIGN (CD209) in the decidua of early human pregnancy. [Am J Pathol. 162: 887-96.](#)
 6. Goddard, S. *et al.* (2004) Interleukin-10 secretion differentiates dendritic cells from human liver and skin. [Am J Pathol. 164: 511-9.](#)
 7. Bournazos, S. *et al.* (2008) Monocyte functional responsiveness after PSGL-1-mediated platelet adhesion is dependent on platelet activation status. [Arterioscler Thromb Vasc Biol. 28: 1491-8.](#)
 8. Hovden, A.O. *et al.* (2011) Maturation of monocyte derived dendritic cells with OK432 boosts IL-12p70 secretion and conveys strong T-cell responses. [BMC Immunol. 12:2.](#)
 9. Hsu, T.L. *et al.* (2002) Modulation of dendritic cell differentiation and maturation by decoy receptor 3. [J Immunol. 168: 4846-53.](#)
 10. Karlsson, H. *et al.* (2002) Innate immune responses of human neonatal cells to bacteria from the normal gastrointestinal flora. [Infect Immun. 70: 6688-96.](#)
 11. Din JN *et al.* (2013) Effect of ω -3 fatty acid supplementation on endothelial function, endogenous fibrinolysis and platelet activation in male cigarette smokers. [Heart. 99 \(3\): 168-74.](#)
 12. Fischer, U. and Koellner, B. (2007) Cross-reactivity of human leukocyte differentiation antigen monoclonal antibodies on carp and rainbow trout cells. [Vet Immunol Immunopathol. 119: 142-55.](#)
 13. Lin, C.W. *et al.* (2005) CD94 1A transcripts characterize lymphoblastic lymphoma/leukemia of immature natural killer cell origin with distinct clinical features. [Blood. 106: 3567-74.](#)
 14. Angel, C.E. *et al.* (2009) Distinctive localization of antigen-presenting cells in human lymph nodes. [Blood. 113: 1257-67.](#)
 15. Chang, Y.C. *et al.* (2004) Modulation of macrophage differentiation and activation by decoy receptor 3. [J Leukoc Biol. 75: 486-94.](#)
 16. Iking-Konert, C. *et al.* (2008) T lymphocytes in patients with primary vasculitis: expansion of CD8+ T cells with the propensity to activate polymorphonuclear neutrophils. [Rheumatology \(Oxford\). 47: 609-16.](#)
 17. Brook, F.A. *et al.* (2010) Derivation and characterisation of the human embryonic stem cell line, OxF1. [In Vitro Cell Dev Biol Anim. 46: 173-7.](#)
 18. Bromberek, J.L. *et al.* (2016) Breed Distribution and Clinical Characteristics of B Cell Chronic Lymphocytic Leukemia in Dogs. [J Vet Intern Med. 30 \(1\): 215-22.](#)
 19. Amouroux, R. *et al.* (2016) *De novo* DNA methylation drives 5hmC accumulation in mouse zygotes. [Nat Cell Biol. 18 \(2\): 225-33.](#)
 20. Spiller, K.L. *et al.* (2016) Differential gene expression in human, murine, and cell line-derived macrophages upon polarization. [Exp Cell Res. 347 \(1\): 1-13.](#)
 21. Kannegieter, N.M. *et al.* (2018) Analysis of NFATc1 amplification in T cells for pharmacodynamic monitoring of tacrolimus in kidney transplant recipients. [PLoS One. 13 \(7\): e0201113.](#)
-

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at:
<https://www.bio-rad-antibodies.com/SDS/MCA596GT>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)
Goat Anti Mouse IgG (STAR76...) [RPE](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
Goat Anti Mouse IgG (STAR77...) [HRP](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

North & South America Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M368543:200529'

Printed on 18 Jan 2024