

# Datasheet: MCA5973F

**BATCH NUMBER 150152**

<b>Description:</b>	MOUSE ANTI PIG CD27:FITC
<b>Specificity:</b>	CD27
<b>Other names:</b>	SWC2
<b>Format:</b>	FITC
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	B30C7
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/10
Immunofluorescence			■	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Pig		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide (NaN <sub>3</sub> )		
Stabilisers	1% Bovine Serum Albumin		

<b>Approx. Protein Concentrations</b>	IgG concentration 0.1 mg/ml
<b>Immunogen</b>	Porcine peripheral blood monocytes.
<b>External Database Links</b>	<b>UniProt:</b> <a href="#">F1SL30</a> <a href="#">Related reagents</a>
<b>Fusion Partners</b>	Spleen cells from immunized Balb/c mice were fused with cells of the SP2/0-Ag14 myeloma cell line
<b>Specificity</b>	<p><b>Mouse anti Pig CD27 antibody, clone B30C7</b> recognizes the porcine homologue of human CD27, previously known as Swine Workshop Cluster 2 (SWC2), a T-cell co-stimulatory molecule belonging to the TNF receptor family. In humans the CD27 antigen is expressed by discrete populations of T- and B-cells where it functions in a co-stimulatory role to induce proliferation of T-cells and B-cells, however studies undertaken using the B30C7 clone have been unable to detect expression of CD27 on porcine B-cells (<a href="#">Reutner et al. 2012</a>).</p> <p>Studies in humans have demonstrated CD27 to be involved in apoptosis, research in swine has shown that porcine CD27 is expressed by all naïve CD8a<sup>-ve</sup> T-helper cells and a sub-population of CD8a<sup>+ve</sup> cells (<a href="#">Reutner et al. 2012</a>).</p> <p>Monoclonal antibodies to CD27 have previously been used to differentiate between subsets of NK cells and research utilizing clone B30C7 has shown that this monoclonal antibody may be used to differentiate between subsets of pig NK cells (<a href="#">Mair et al. 2013</a>).</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul
<b>References</b>	<ol style="list-style-type: none"> <li>1. Reutner, K. <i>et al.</i> (2012) Porcine CD27: identification, expression and functional aspects in lymphocyte subsets in swine. <a href="#">Dev Comp Immunol. 38: 321-31.</a></li> <li>2. Reutner, K. <i>et al.</i> (2013) CD27 expression discriminates porcine T helper cells with functionally distinct properties. <a href="#">Vet Res. 44: 18.</a></li> <li>3. Franzoni, G. <i>et al.</i> (2013) Assessment of the Phenotype and Functionality of Porcine CD8 T Cell Responses following Vaccination with Live Attenuated Classical Swine Fever Virus (CSFV) and Virulent CSFV Challenge. <a href="#">Clin Vaccine Immunol. 20: 1604-16.</a></li> <li>4. Mair, K.H. <i>et al.</i> (2013) Porcine CD8αdim/-NKp46high NK cells are in a highly activated state. <a href="#">Vet Res. 44: 13.</a></li> <li>5. López, E. <i>et al.</i> (2019) Identification of very early inflammatory markers in a porcine myocardial infarction model. <a href="#">BMC Vet Res. 15 (1): 91.</a></li> </ol>
<b>Storage</b>	<p>Store at +4°C or at -20°C if preferred.</p> <p>This product should be stored undiluted.</p> <p>Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.</p>

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

<b>Guarantee</b>	12 months from date of despatch
------------------	---------------------------------

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10041 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA5973F">https://www.bio-rad-antibodies.com/SDS/MCA5973F</a> 10041
--------------------------------------	--

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)  
'M368554:200529'

**Printed on 28 May 2025**