

Datasheet: MCA5973F

BATCH NUMBER 166716

Description:	MOUSE ANTI PIG CD27:FITC
Specificity:	CD27
Other names:	SWC2
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	B30C7
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10
Immunofluorescence			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Pig		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide (NaN ₃)		
Stabilisers	1% Bovine Serum Albumin		

Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	Porcine peripheral blood monocytes.
External Database Links	UniProt: F1SL30 Related reagents
Fusion Partners	Spleen cells from immunized Balb/c mice were fused with cells of the SP2/0-Ag14 myeloma cell line
Specificity	<p>Mouse anti Pig CD27 antibody, clone B30C7 recognizes the porcine homologue of human CD27, previously known as Swine Workshop Cluster 2 (SWC2), a T-cell co-stimulatory molecule belonging to the TNF receptor family. In humans the CD27 antigen is expressed by discrete populations of T- and B-cells where it functions in a co-stimulatory role to induce proliferation of T-cells and B-cells, however, using the B30C7 clone, expression of CD27 on porcine B-cells appears undetectable (Reutner et al. 2012).</p> <p>Porcine CD27 is expressed by all naïve CD8a^{ve} T-helper cells and a sub-population of CD8a^{ve} cells (Reutner et al. 2012).</p> <p>Monoclonal antibodies to CD27 have previously been used to differentiate between subsets of NK cells and clone B30C7 may be used to differentiate between subsets of pig NK cells (Mair et al. 2013).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul
References	<ol style="list-style-type: none"> 1. Reutner, K. <i>et al.</i> (2012) Porcine CD27: identification, expression and functional aspects in lymphocyte subsets in swine. Dev Comp Immunol. 38: 321-31. 2. Reutner, K. <i>et al.</i> (2013) CD27 expression discriminates porcine T helper cells with functionally distinct properties. Vet Res. 44: 18. 3. Mair, K.H. <i>et al.</i> (2013) Porcine CD8αdim/-NKp46high NK cells are in a highly activated state. Vet Res. 44: 13. 4. Franzoni, G. <i>et al.</i> (2013) Assessment of the Phenotype and Functionality of Porcine CD8 T Cell Responses following Vaccination with Live Attenuated Classical Swine Fever Virus (CSFV) and Virulent CSFV Challenge. Clin Vaccine Immunol. 20: 1604-16. 5. López, E. <i>et al.</i> (2019) Identification of very early inflammatory markers in a porcine myocardial infarction model. BMC Vet Res. 15 (1): 91. 6. Maciag, S.S. <i>et al.</i> (2022) The influence of source of porcine colostrum in development of early immune ontogeny in the piglet Res Sq. Mar 24 [Epub ahead of print]. 7. Bettin, L. <i>et al.</i> (2023) Co-stimulation by TLR7/8 ligand R848 modulates IFN-γ production of porcine γδ T cells in a microenvironment-dependent manner. Dev Comp Immunol. 138: 104543. 8. Haach, V. <i>et al.</i> (2023) A polyvalent virosomal influenza vaccine induces broad cellular and humoral immunity in pigs. Virol J. 20 (1): 181.
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at

-20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA5973F 10041
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M384685:210513'

Printed on 18 Jan 2024