

Datasheet: MCA5974F

BATCH NUMBER 170104

Description:	MOUSE ANTI PIG CD52:FITC
Specificity:	CD52
Other names:	SWC1
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	11/305/44
Isotype:	IgG2b
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10
Immunofluorescence			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Pig		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide (NaN ₃)		
Stabilisers	1% Bovine Serum Albumin		

Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	Porcine thymocytes
External Database Links	UniProt: H8ZRT6 Related reagents
Fusion Partners	Spleen cells from immunized Balb/c mice were fused with cells of the SP2/0-Ag14 myeloma cell line
Specificity	<p>Mouse anti Pig CD52, clone 11/305/44 recognizes the porcine homologue of human CD52, a ~19 kDa antigen expressed by mature lymphocytes, monocytes and dendritic cells.</p> <p>Mouse anti Pig CD52, clone 11/305/44 was originally clustered at the 1st International Swine Cluster of Differentiation Workshop held in 1992 as SWC1 (Lunney et al. 1994). SWC1 is the porcine orthologue to human CD52, expressed by most leucocytes including resting T-cells, monocytes and granulocytes, but is not expressed by B-cells, erythrocytes or platelets (Piriou-Guyzlack et al. 2008) & (Leitner et al. 2012).</p> <p>Porcine CD52, expressed at very much higher levels on monocytes than mature macrophages, and SWC9, expressed exclusively on mature tissue macrophages, have been used as markers of monocyte-macrophage differentiation (Sanchez et al. 1999) & (McCullough et al. 1999).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul
References	<ol style="list-style-type: none"> Lunney, J.K. <i>et al.</i> (1994) Overview of the First International Workshop to Define Swine Leukocyte Cluster of Differentiation (CD) Antigens. Vet Immunol Immunopathol. 43: 193-206. Leitner, J. <i>et al.</i> (2012) Porcine SWC1 is CD52--final determination by the use of a retroviral cDNA expression library. Vet Immunol Immunopathol. 146 (1): 27-34. Seeboth, J. <i>et al.</i> (2012) The fungal T-2 toxin alters the activation of primary macrophages induced by TLR-agonists resulting in a decrease of the inflammatory response in the pig. Vet Res. 43: 35. Shao, L. <i>et al.</i> (2016) Tissue-specific mRNA expression profiles of porcine Toll-like receptors at different ages in germ-free and conventional pigs. Vet Immunol Immunopathol. 171: 7-16.
Further Reading	<ol style="list-style-type: none"> Sánchez, C. <i>et al.</i> (1999) The Porcine 2A10 Antigen is Homologous to Human CD163 and Related to Macrophage Differentiation. J Immunol. 162: 5230-7. McCullough, K.C. <i>et al.</i> (1999) Intermediate stages in monocyte-macrophage differentiation modulate phenotype and susceptibility to virus infection. Immunology. 98 (2): 203-12. Piriou-Guyzlack, L. <i>et al.</i> (2008) Membrane markers of the immune cells in swine: an update. Vet Res. 39: 54

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA5974F>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG2b NEGATIVE CONTROL:FITC \(MCA691F\)](#)

Recommended Useful Reagents

[MOUSE ANTI PIG CD203a \(MCA1973GA\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M384623:210513'

Printed on 29 Jan 2026