

Datasheet: MCA597GA

BATCH NUMBER 0814

Description:	MOUSE ANTI HUMAN COLLAGEN VII
Specificity:	COLLAGEN VII
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	LH7.2
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen	▪			1/25 - 1/100
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	Does not react with:Rat, Mouse, Pig
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide

Stabilisers

Carrier Free Yes

Approx. Protein Concentrations IgG concentration 1.0 mg/ml

Immunogen Crude extract of skin

External Database Links

UniProt:

[Q02388](#)

[Related reagents](#)

Entrez Gene:

[1294](#)

COL7A1

[Related reagents](#)

Fusion Partners Spleen cells from immunised BALB/c mice were fused with cells of the mouse Sp2 myeloma cell line

Specificity

Mouse anti Human collagen VII antibody, clone LH7.2 recognizes the carboxy-terminus of type VII collagen ([Kirkham *et al.* 1989](#)). Collagen VII is a basement membrane protein in stratified squamous epithelia which is involved in membrane organization through interaction with other ECM components ([Gammon *et al.* 1992](#)). Collagen VII is composed of three identical alpha chains, each having an amino-terminal non-collagenous domain and a carboxy-terminal collagenous domain.

Mouse anti Human collagen VII, clone LH7.2 has proved effective for immunohistochemical identification of basement membrane in human skin ([Watson *et al.* 2001](#)).

References

1. Heagerty, A.H. *et al.* (1986) Identification of an epidermal basement membrane defect in recessive forms of dystrophic epidermolysis bullosa by LH 7:2 monoclonal antibody: use in diagnosis. [Br J Dermatol. 115 \(2\): 125-31.](#)
2. Leigh, I.M. *et al.* (1987) LH7.2 Monoclonal antibody detects type VII collagen in the sublamina densa zone of ectodermally-derived epithelia, including skin. *Epithelia* 1: 17-29.
3. Kirkham, N. *et al.* (1989) Type VII collagen antibody LH 7.2 identifies basement membrane characteristics of thin malignant melanomas. [J Pathol. 157 \(3\): 243-7.](#)
4. Leigh, I.M. *et al.* (1988) Type VII collagen is a normal component of epidermal basement membrane, which shows altered expression in recessive dystrophic epidermolysis bullosa. [J Invest Dermatol. 90 \(5\): 639-42.](#)
5. Ghohestani, R.F. *et al.* (1998) IgE antibodies in sera from patients with bullous pemphigoid are autoantibodies preferentially directed against the 230-kDa epidermal antigen (BP230). [J Clin Immunol. 18: 202-9.](#)
6. Craven, N.M. *et al.* (1997) Clinical features of photodamaged human skin are associated with a reduction in collagen VII. [Br J Dermatol. 137: 344-50.](#)
7. Al-Refu, K and Goodfield, M. (2011) Immunohistochemistry of ultrastructural changes in scarring lupus erythematosus. [Clin Exp Dermatol. 36: 63-8.](#)
8. Herndon, D.N. *et al.* (1995) Characterization of growth hormone enhanced donor site

healing in patients with large cutaneous burns. [Ann Surg. 221: 649-56.](#)

9. Al-Refu, K. and Goodfield, M. (2009) Basement membrane changes in lichen planopilaris. [J Eur Acad Dermatol Venereol. 23: 1289-93.](#)

10. Watson, R.E. *et al.* (2001) A short-term screening protocol, using fibrillin-1 as a reporter molecule, for photoaging repair agents. [J Invest Dermatol. 116: 672-8.](#)

11. Jolicoeur, F. *et al.* (2003) Basal cells of second trimester fetal breasts: immunohistochemical study of myoepithelial precursors. [Pediatr Dev Pathol. 6: 398-413.](#)

Storage

Store at +4°C for one month or at -20°C for longer.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA597GA>
10040

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP

North & South America	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M368561:200529'

Printed on 18 Jan 2024