# Datasheet: MCA5983GA BATCH NUMBER 1114

Description:	MOUSE ANTI HORSE IgE
Specificity:	lgE
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	7H2
lsotype:	lgG1
Quantity:	0.1 mg

# **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .				
		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry			•	
	Immunohistology - Frozen	-			
	Immunohistology - Paraffin (1)	•			
	ELISA	-			1/40 - 1/400
	Immunoprecipitation				
	Western Blotting (2)	-			
	necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls. (1) This antibody is demonstrated to work in formalin fixed tissue when using trypsin digestion pre-treatment of paraffin sections. Testing undertaken utilizing heat mediated antigen retrieval techniques with this antibody have not been successful. (2) Non-reducing conditions are recommended				
Target Species	Horse				
Product Form	Purified IgG - liquid				
Preparation	Purified IgG prepared by supernatant	affinity c	hromatogi	aphy on Protein A fror	m tissue culture

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN <sub>3</sub> )
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Equine IgE
Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of the P3X myeloma cell line
Specificity	Mouse anti Horse IgE antibody, clone 7H2 recognizes native equine IgE and does not cross react with equine IgM, IgA or IgG.
	IgE is an immunoglobulin primarily produced from plasma cells and, in normal serum, present at very low concentrations. Western blot analysis against affinity purified equine IgE using Mouse anti Horse IgE clone 7H2 demonstrates a single major band of approximately 200 kDa under non-reducing conditions that corresponds with the expected molecular weight of the complete equine IgE molecule ( <u>Wilson, D.A. <i>et al.</i> 2006</u> ). IgE is important in both type 1 hypersensitivity and immunity to parasite infections, in
	particular parasitic worms where equine IgE levels are significantly elevated following worm infection.
	Monoclonal antibodies to equine IgE are of particular relevance to research into insect bite sensitivity, one of the most widely studied allergic diseases in equid species ( <u>Schaffartzik,</u> <u>A., <i>et al.</i> 2012</u> ).
ELISA	This product may be used in an indirect ELISA or as a capture antibody in a sandwich ELISA together with MCA5982P as the detection reagent.
Western Blotting	Clone 7H2 has been demonstrated to detect a band of approximately 200 kDa in non-reducing conditions. This clone is not suitable for use in western blotting under reducing conditions. For western blotting applications it is recommended to use clone 3H10 <u>MCA5982GA</u> .
References	1. Wilson, D. A. <i>et al.</i> (2006) Production of monoclonal antibodies specific for native equine IgE and their application to monitor total serum IgE responses in Icelandic and non-Icelandic horses with insect bite dermal hypersensitivity. <u>Vet Immunol Immunopathol.</u> <u>112: 156-70.</u>
Further Reading	1. Schaffartzik, A. <i>et al.</i> (2012) Equine insect bite hypersensitivity: what do we know? <u>Vet</u> Immunol Immunopathol. 147: 113-26
Storage	Store at +4°C or at -20°C if preferred.

	Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.	,
Guarantee	12 months from date of despatch	
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA5983GA 10040	
Regulatory	For research purposes only	

## **Related Products**

### **Recommended Secondary Antibodies**

Rabbit A	nti Mouse IgG (STAR12)	RPE	<u> </u>		
Goat Ant	i Mouse IgG IgA IgM (STAR87	) <u>HR</u> F	2		
Goat Ant	i Mouse IgG (STAR76)	RPE			
Rabbit A	nti Mouse IgG (STAR13)	HR	-		
Goat Ant	i Mouse IgG (STAR70)	<u>FIT(</u>	<u>C</u>		
Goat Ant	i Mouse IgG (H/L) (STAR117)	<u>Alk.</u>	Phos., DyLight®488, Dy	<u>_ight®550,</u>	
		DyL	ight®650, DyLight®680,	DyLight®800	),
		FIT(	<u>C, HRP</u>		
Rabbit A	nti Mouse IgG (STAR9)	<u>FIT(</u>	<u>C</u>		
Goat Ant	i Mouse IgG (STAR77)	HR	-		
Goat Ant	i Mouse IgG (Fc) (STAR120)	<u>FIT(</u>	<u>C, HRP</u>		
Recomm	nended Useful Reagents				
MOUSE A	NTI HORSE IgGb (MCA1901GA) NTI HORSE IgE (MCA5982GA) NTI HORSE IgE:HRP (MCA5982P)				
North & South	Tel: +1 800 265 7376 Worldwig	de	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com		Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rac	.com	Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets M368566:200529'

#### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint