### Datasheet: MCA5985GA BATCH NUMBER 155948

Description:	MOUSE ANTI HUMAN IP-10
Specificity:	IP-10
Other names:	CXCL10
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	A02-1E10
Isotype:	lgG2b
Quantity:	0.1 mg

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
		Yes	No	Not Determined	Suggested Dilution	
	ELISA					
	Where this product has no necessarily exclude its us a guide only. It is recomm system using appropriate	e in such ended the	procedur at the use	es. Suggested workir r titrates the product	ng dilutions are given as	
Target Species	Human					
Product Form	Purified IgG - liquid					
Preparation	Purified IgG prepared by affinity chromatography on Protein G from serum free tissue culture supernatant					
Buffer Solution	Phosphate buffered saline					
Preservative Stabilisers	0.09% Sodium Azide (NaN <sub>3</sub> )					
Approx. Protein Concentrations	IgG concentration 1.0 mg	/ml				
Immunogen	<i>E. coli</i> derived recombina	nt protein	correspo	nding to aa 22-98 of	human IP-10	

External Database Links	UniProt: <u>P02778</u> Related reagents Entrez Gene:			
	3627 CXCL10 Related reagents			
Synonyms	INP10, SCYB10			
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the SP2/0 myeloma cell line			
Specificity	<b>Mouse anti Human IP-10, clone A02-1E10</b> recognizes the chemokine IP-10. IP-10, also known as CXCL10, is a small cytokine belonging to the CXC chemokine family. IP-10 cDNA encodes a 98 amino acid protein which is cleaved and secreted from a wide variety of cell types in response to interferon gamma and other inflammatory cytokines ( <u>Ohmori &amp; Hamilton 1995</u> ).			
	IP-10 exerts its biological effects mainly via binding to CXCR3 ( <u>van den Borne <i>et al.</i></u> <u>2014</u> ). Also, CXCR3-independent pathways have been studied for IP-10/CXCL10 involving binding to heparan sulfate glycosaminoglycans (GAGs). IP-10 not only induces chemoattraction of inflammatory cells, but also migration and proliferation of endothelial cells and VSMCs ( <u>Wang <i>et al.</i> 1996</u> ).			
Further Reading	<ol> <li>van den Borne, P. <i>et al.</i> (2014) The multifaceted functions of CXCL10 in cardiovascular disease <u>Biomed Res Int. 2014: 893106.</u></li> <li>Ohmori, Y. &amp; Hamilton, T.A. (1994) Cell type and stimulus specific regulation of chemokine gene expression <u>Biochem Biophys Res Commun. 198: 590-6.</u></li> <li>Wang, X. <i>et al.</i> (1996) Interferon-inducible protein-10 involves vascular smooth muscle cell migration, proliferation, and inflammatory response <u>J Biol Chem. 271: 24286-93.</u></li> </ol>			
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.			
Guarantee	12 months from date of despatch			
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA5985GA 10040			
Regulatory	For research purposes only			

# **Related Products**

## **Recommended Secondary Antibodies**

Rabbit Ar	nti Mouse IgG (STAR12)	RPE				
	i Mouse IgG IgA IgM (STAR87					
Goat Anti	Mouse IgG (STAR76)	RPE				
Rabbit Ar	nti Mouse IgG (STAR13)	HRP				
Goat Ant	i Mouse IgG (STAR70)	FITC				
Goat Anti	i Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLight®550,				
		DyLight®650, DyLight®680, DyLight®800	<u>0,</u>			
		<u>FITC, HRP</u>				
Rabbit Ar	nti Mouse IgG (STAR9)	FITC				
Goat Ant	i Mouse IgG (STAR77)	HRP				
Goat Ant	Mouse IgG (Fc) (STAR120)	<u>FITC, HRP</u>				
North & South	Tel: +1 800 265 7376 Worldwid	(1) 11 11 11 11 11	Tel: +49 (0) 89 8090 95 21			
America	Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com			

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M368569:200529'

#### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint