

## Datasheet: MCA5997

<b>Description:</b>	RAT ANTI MOUSE CD64
<b>Specificity:</b>	CD64
<b>Other names:</b>	FcRI
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	AT152-9
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	0.25 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10
Immunohistology - Frozen	▪			1/1000

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by ion exchange chromatography from tissue culture supernatant
<b>Buffer Solution</b>	TRIS buffered saline
<b>Preservative Stabilisers</b>	<0.1% Sodium Azide (NaN <sub>3</sub> )
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	HEK293F cells expressing CD64 and gamma chain

**External Database  
Links**

**UniProt:**

[P26151](#)   [Related reagents](#)

**Entrez Gene:**

[14129](#) Fcgr1   [Related reagents](#)

---

**Synonyms**

Fcg1

---

**RRID**

AB\_2687456

---

**Fusion Partners**

Spleen cells from immunised rats were fused with cells of the NS-1 myeloma cell line

---

**Specificity**

**Rat anti mouse CD64 antibody, clone AT152-9**, recognizes mouse CD64 also known as FcRI. CD64 is a high affinity activatory receptor for IgG2a and a low affinity receptor for IgG2b and IgG3 type antibodies. The interaction between Fc receptors and antibodies play important roles in both the innate and adaptive immune responses.

CD64, through binding of the Fc segment of IgG, mediates phagocytosis and plays a role in antibody-dependent cellular cytotoxicity and clearance of immune complexes. In addition, CD64 also functions as an antigen capture for presentation to T-cells and also mediates the release of cytokines and reactive oxygen intermediates including interleukin (IL)-1, IL-6 and tumor necrosis factor (TNF) alpha. It is constitutively expressed on monocytes and macrophages, germinal centre dendritic cells and early myeloid lineage cells, but not lymphocytes. Expression on monocytes can be strongly upregulated by treatment with interferon (IFN) gamma or G-CSF, and can be induced on neutrophils and eosinophils by IFN gamma.

---

**Flow Cytometry**

Use 10ul of the suggested working dilution to label  $1 \times 10^6$  cells in 100ul

---

**References**

1. White; A.L. *et al.* (2011) Interaction with FcγRIIB is critical for the agonistic activity of anti-CD40 monoclonal antibody. [J Immunol. 187 \(4\): 1754-63.](#)
2. Tutt AL *et al.* (2015) Development and Characterization of Monoclonal Antibodies Specific for Mouse and Human Fcγ Receptors. [J Immunol. 195 \(11\): 5503-16.](#)

---

**Storage**

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

**Guarantee**

12 months from date of despatch

---

**Health And Safety  
Information**

Material Safety Datasheet documentation #10057 available at:  
<https://www.bio-rad-antibodies.com/SDS/MCA5997>  
10057

---

**Regulatory**

For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Goat Anti Rat IgG (STAR69...)	<a href="#">FITC</a>
Goat Anti Rat IgG (STAR73...)	<a href="#">RPE</a>
Rabbit Anti Rat IgG (STAR16...)	<a href="#">DyLight@800</a>
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	<a href="#">DyLight@650</a> , <a href="#">DyLight@800</a>
Goat Anti Rat IgG (STAR72...)	<a href="#">HRP</a>
Rabbit Anti Rat IgG (STAR21...)	<a href="#">HRP</a>
Rabbit Anti Rat IgG (STAR17...)	<a href="#">FITC</a>
Goat Anti Rat IgG (STAR131...)	<a href="#">Alk. Phos.</a> , <a href="#">Biotin</a>

### Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL \(MCA1212\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------	------------------	-------------------------------------------------------------------------------------------------------------------------------------------------	---------------	-----------------------------------------------------------------------------------------------------------------------------------------------------

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)  
'M389649:210806'

Printed on 25 Mar 2023