Datasheet: MCA5997F BATCH NUMBER 148818

| Description: | RAT ANTI MOUSE CD64:FITC | | | |
|---------------|--------------------------|--|--|--|
| Specificity: | CD64 | | | |
| Other names: | FcRI | | | |
| Format: | FITC | | | |
| Product Type: | Monoclonal Antibody | | | |
| Clone: | AT152-9 | | | |
| lsotype: | lgG2a | | | |
| Quantity: | 0.1 mg | | | |

Product Details

| Applications | This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u> rad-antibodies.com/protocols. | | | | | | | |
|-----------------------------------|--|-----------------------------------|--------------------|---|--------------------------|--|--|--|
| | | Yes | No | Not Determined | Suggested Dilution | | | |
| | Flow Cytometry | • | | | Neat | | | |
| | Where this product has necessarily exclude its a guide only. It is recor system using appropria | use in such pro mmended that t | ocedure he user | s. Suggested workin titrates the product f | g dilutions are given as | | | |
| Target Species | Mouse | | | | | | | |
| Product Form | Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid | | | | | | | |
| Max Ex/Em | Fluorophore | Excitation Max | (nm) E | Emission Max (nm) | | | | |
| | FITC | 490 | | 525 | | | | |
| Preparation | Purified IgG prepared by ion exchange chromatography from tissue culture supernatant | | | | | | | |
| Buffer Solution | Phosphate buffered saline | | | | | | | |
| Preservative Stabilisers | 0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin | | | | | | | |
| Approx. Protein Concentrations | IgG concentration 0.1 | mg/ml | | | | | | |

| External Database Links | UniProt: <u>P26151</u> <u>Related reagents</u> Entrez Gene: <u>14129</u> Fcgr1 <u>Related reagents</u> | | | | |
|----------------------------------|---|--|--|--|--|
| | <u>14129</u> i cgi i <u>Related reagents</u> | | | | |
| Synonyms | Fcg1 | | | | |
| Fusion Partners | Spleen cells from immunised rats were fused with cells of the NS-1 myeloma cell line | | | | |
| Specificity | Rat anti mouse CD64 antibody, clone AT152-9 , recognizes mouse CD64 also known as FcRI. CD64 is a high affinity activatory receptor for IgG2a and a low affinity receptor for IgG2b and IgG3 type antibodies. The interaction between Fc receptors and antibodies play important roles in both the innate and adaptive immune responses. | | | | |
| | CD64, through binding of the Fc segment of IgG, mediates phagocytosis and plays a role in antibody-dependent cellular cytotoxicity and clearance of immune complexes. In addition, CD64 also functions as an antigen capture for presentation to T-cells and also mediates the release of cytokines and reactive oxygen intermediates including interleukin (IL)-1, IL-6 and tumor necrosis factor (TNF) alpha. It is constitutively expressed on monocytes and macrophages, germinal centre dendritic cells and early myeloid lineage cells, but not lymphocytes. Expression on monocytes can be strongly upregulated by treatment with interferon (IFN) gamma or G-CSF, and can be induced on neutrophils and eosinophils by IFN gamma. | | | | |
| Flow Cytometry | Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul | | | | |
| References | White; A.L. <i>et al.</i> (2011) Interaction with FcγRIIB is critical for the agonistic activity of anti-CD40 monoclonal antibody. <u>J Immunol. 187 (4): 1754-63.</u> Tutt AL <i>et al.</i> (2015) Development and Characterization of Monoclonal Antibodies Specific for Mouse and Human Fcγ Receptors. <u>J Immunol. 195 (11): 5503-16.</u> | | | | |
| Storage | Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. | | | | |
| Guarantee | 12 months from date of despatch | | | | |
| Health And Safety Information | Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA5997F 10041 | | | | |
| Regulatory | For research purposes only | | | | |
| | | | | | |

Related Products

Recommended Negative Controls

RAT IgG2a NEGATIVE CONTROL:FITC (MCA1212F)

| North & South | Tel: +1 800 265 7376 | Worldwide | Tel: +44 (0)1865 852 700 | Europe | Tel: +49 (0) 89 8090 95 21 |
|---------------|---------------------------------|-----------|---------------------------------|--------|--------------------------------------|
| America | Fax: +1 919 878 3751 | | Fax: +44 (0)1865 852 739 | | Fax: +49 (0) 89 8090 95 50 |
| | Email: antibody_sales_us@bio-ra | ad.com | Email: antibody_sales_uk@bio-ra | ad.com | Email: antibody_sales_de@bio-rad.com |

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M368577:200529'

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint