

Datasheet: MCA5998

Description:	RAT ANTI MOUSE CD16
Specificity:	CD16
Other names:	FcRIII
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	AT154-2
Isotype:	IgG2b
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10
Immunohistology - Frozen	▪			1/100 - 1/1000

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	TRIS buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	HEK293F cell expressing CD16 and gamma chain

**External Database
Links**

UniProt:

[P08508](#) [Related reagents](#)

Entrez Gene:

[14131](#) Fcgr3 [Related reagents](#)

Fusion Partners Spleen cells from immunised rats were fused with cells of the NS-1 myeloma cell line

Specificity **Rat anti Mouse CD16 antibody, clone AT154-2**, recognizes CD16, also known as FcR111. CD16 is a low affinity activatory receptor which interacts preferentially with mouse IgG1, IgG2a and IgG2b immune complexes and is involved in various effector functions including phagocytosis, NK antibody dependent cellular cytotoxicity and the release of inflammatory mediators.

In mice, CD16 is expressed on macrophages, NK cells, granulocytes, myeloid precursors and subpopulations of T lymphocytes.

Flow Cytometry Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.

References

1. White AL *et al.* (2011) Interaction with FcγRIIB is critical for the agonistic activity of anti-CD40 monoclonal antibody. [J Immunol. 187 \(4\): 1754-63.](#)
2. Tutt, A.L. *et al.* (2015) Development and Characterization of Monoclonal Antibodies Specific for Mouse and Human Fcγ Receptors. [J Immunol. 195 \(11\): 5503-16.](#)

Storage

Store at +4°C or at -20°C if preferred.
Storage in frost-free freezers is not recommended.
This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10057 available at: 10057: <https://www.bio-rad-antibodies.com/uploads/MSDS/10057.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Rabbit Anti Rat IgG (STAR16...)	DyLight®800

Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) [DyLight®800](#)

Goat Anti Rat IgG (STAR72...) [HRP](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M368578:200529'

Printed on 22 Mar 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)