

Datasheet: MCA6004AMO

Description:	RAT IgG1 NEGATIVE CONTROL:Amethyst Orange
Specificity:	RAT IgG1 NEGATIVE CONTROL
Format:	Amethyst Orange
Product Type:	Negative/Isotype Control
Isotype:	lgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			*

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. *This antibody should be used at the same concentration as the test antibody.

Target Species	Negative Control	Negative Control Purified IgG conjugated to Amethyst Orange - liquid				
Product Form	Purified IgG conjuga					
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm	n)		
	Amethyst Orange	405	540			
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant					
Buffer Solution	Phosphate buffered					
Preservative Stabilisers	0.09% Sodium Azid 1% Bovine Serum A					
Approx. Protein Concentrations	IgG concentration 0	.1 mg/ml				
Immunogen	Horseradish peroxic	lase				
Specificity	Rat IgG1 negative	control, a rat monoclor	al raised against h	orseradish peroxidase		

Test results have shown that this antibody is also suitable for use as a negative control with porcine and canine cells. Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul References 1. do Prado Duzanski, A. et al. (2022) Cell-mediated immunity and expression of MHC class I and class II molecules in dogs naturally infected by canine transmissible venereal tumor: Is there complete spontaneous regression outside the experimental CTVT? Research in Veterinary Science. 145: 193-204. 2. Matralis, D.T. et al. (2023) Intracellular IFN-γ and IL-4 levels of CD4 + and CD8 + T cells in the peripheral blood of naturally infected (Leishmania infantum) symptomatic dogs before and following a 4-week treatment with miltefosine and allopurinol: a double-blinded, controlled and cross-sectional study. Acta Vet Scand. 65 (1): 2. **Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light. Guarantee 12 months from date of despatch **Health And Safety** Material Safety Datasheet documentation #10041 available at: Information https://www.bio-rad-antibodies.com/SDS/MCA6004AMO 10041 Regulatory For research purposes only North & South Tel: +1 800 265 7376 Tel: +44 (0)1865 852 700 Worldwide Europe Tel: +49 (0) 89 8090 95 21

recommended for use as a negative control to assess the level of non-specific binding of rat IgG1 test antibodies to the surface of human and mouse cells in flow cytometry.

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M395453:220429'

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

America