Datasheet: MCA6004EL

BATCH NUMBER 166477

Description:	RAT IgG1 NEGATIVE CONTROL:Low Endotoxin
Specificity:	RAT IgG1 NEGATIVE CONTROL
Format:	Low Endotoxin
Product Type:	Negative/Isotype Control
Isotype:	IgG1
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			*

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. *This antibody should be used at the same concentration as the test antibody.

Target Species	Negative Control
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	None present
Endotoxin Level	<0.01EU/ug
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Horseradish peroxidase
Specificity	**Rat IgG1 negative control**, a rat monoclonal raised against horseradish peroxidase, is

	recommended for use as a negative control to assess the level of non-specific binding of rat IgG1 test antibodies to the surface of human and mouse cells in flow cytometry. Test results have shown that this antibody is also suitable for use as a negative control with porcine and canine cells.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10^6 cells in 100ul
References	1. do Prado Duzanski, A. *et al.* (2022) Cell-mediated immunity and expression of MHC class I and class II molecules in dogs naturally infected by canine transmissible venereal tumor: Is there complete spontaneous regression outside the experimental CTVT? Research in Veterinary Science. 145: 193-204. 2. Matralis, D.T. *et al.* (2023) Intracellular IFN-γ and IL-4 levels of CD4 + and CD8 + T cells in the peripheral blood of naturally infected (*Leishmania infantum*) symptomatic dogs before and following a 4-week treatment with miltefosine and allopurinol: a double-blinded, controlled and cross-sectional study. Acta Vet Scand. 65 (1): 2.
Storage	Store at -20°C only. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/MCA6004EL 10162
Regulatory	For research purposes only

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	**Worldwide**	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	**Europe**	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M395455:220429'

Printed on 28 Aug 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint