

Datasheet: MCA6005F

BATCH NUMBER 169176

Description:	RAT IgG2a NEGATIVE CONTROL:FITC
Specificity:	RAT IgG2a NEGATIVE CONTROL
Format:	FITC
Product Type:	Negative/Isotype Control
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			*

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. *This antibody should be used at the same concentration as the test antibody.

Target Species	Negative Control		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide (NaN ₃)		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml		
Immunogen	Trinitrophenol		

Specificity **Rat IgG2a negative control**, a rat monoclonal raised against trinitrophenol, is recommended for use as a negative control to assess the level of non-specific binding of rat IgG2a test antibodies to the surface of human and mouse cells in flow cytometry.

Test results have shown that this antibody is also suitable for use as a negative control with porcine and canine cells.

Flow Cytometry Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul

References

1. Barraud-Lange, V. *et al.* (2020) Partial Sperm beta1 Integrin Subunit Deletion Proves its Involvement in Mouse Gamete Adhesion/Fusion. [Int J Mol Sci. 21 \(22\)Nov 11 \[Epub ahead of print\].](#)
2. do Prado Duzanski, A. *et al.* (2022) Cell-mediated immunity and expression of MHC class I and class II molecules in dogs naturally infected by canine transmissible venereal tumor: Is there complete spontaneous regression outside the experimental CTVT? [Research in Veterinary Science. 145: 193-204.](#)
3. Karayannopoulou, M. *et al.* (2022) Effect of major versus minor mastectomy on host immunity in canine mammary cancer [Vet Immunol Immunopathol. Feb 24: 110403.](#)
4. Matralis, D.T. *et al.* (2023) Intracellular IFN- γ and IL-4 levels of CD4 + and CD8 + T cells in the peripheral blood of naturally infected (*Leishmania infantum*) symptomatic dogs before and following a 4-week treatment with miltefosine and allopurinol: a double-blinded, controlled and cross-sectional study. [Acta Vet Scand. 65 \(1\): 2.](#)
5. Rogato, F. *et al.* (2024) Leukemia cutis as a prominent clinical sign in a dog with acute myeloid leukemia. [Vet Clin Pathol. 53 \(4\): 448-57.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at $2-8^{\circ}\text{C}$ for short term use (up to 4 weeks) and store the remaining aliquots at -20°C .

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA6005F>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:FITC \(MCA1212F\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

