

## Datasheet: MCA6005PECY7 BATCH NUMBER 166057

Description:	RAT IgG2a NEGATIVE CONTROL:RPE-Cy7		
Specificity:	RAT IgG2a NEGATIVE CONTROL		
Format:	RPE-CY7		
Product Type:	Negative/Isotype Control		
Isotype:	lgG2a		
Quantity:	100 TESTS/1ml		

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .						
	rad-antipodies.com/pr		No	Not Determined	Suggested Dilution		
	Flow Cytometry	•			*		
	Where this antibody h	as not been test	ted for	use in a particular tec	hnique this does not		
	necessarily exclude its use in such procedures. *This antibody should be used at the						
	same concentration a	s the test antibo	dy.				
Target Species	Negative Control						
Product Form	Purified IgG conjugate	ed to R. Phycoer	ythrin	(RPE)-Cy7 - lyophilize	ed		
Reconstitution	Reconstitute with 1.0 ml distilled water						
Max Ex/Em	Fluorophore	Excitation Max	(nm)	Emission Max (nm)			
	RPE-Cy7 488nm laser	496		695			
Preparation	Purified IgG prepared supernatant	by affinity chron	natogra	aphy on Protein G fror	n tissue culture		
Buffer Solution	Phosphate buffered sa	aline					
Preservative	0.09% Sodium Azide	(NaN <sub>3</sub> )					
Stabilisers	1% Bovine Serum Alb						
	5% Sucrose						
Immunogen	Trinitrophenol						

Regulator th & South erica	Tel: +1 800 265 Fax: +1 919 878	
Informatio		Material Safety Datasheet documentation #20487 available at: <u>https://www.bio-rad-antibodies.com/SDS/MCA6005PECY7</u> 20487
Acknowle	edgements	Cy® and Cydye® are registered trademarks of GE Healthcare
Guarante	e	12 months from date of despatch
Storage		Prior to reconstitution store at +4°C. After reconstitution store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosens and should be protected from light.
Celerence		<ol> <li>Bahadd-Lange, V. <i>et al.</i> (2020) Partial Sperin beta Fintegrin Subunit Deletion Prove Involvement in Mouse Gamete Adhesion/Fusion. <u>Int J Mol Sci. 21 (22)Nov 11 [Epub</u> <u>ahead of print].</u></li> <li>do Prado Duzanski, A. <i>et al.</i> (2022) Cell-mediated immunity and expression of MHC class I and class II molecules in dogs naturally infected by canine transmissible venere tumor: Is there complete spontaneous regression outside the experimental CTVT? <u>Research in Veterinary Science. 145: 193-204.</u></li> <li>Karayannopoulou, M. <i>et al.</i> (2022) Effect of major versus minor mastectomy on hos immunity in canine mammary cancer <u>Vet Immunol Immunopathol. Feb 24: 110403.</u></li> <li>Matralis, D.T. <i>et al.</i> (2023) Intracellular IFN-γ and IL-4 levels of CD4 + and CD8 + T of in the peripheral blood of naturally infected (<i>Leishmania infantum</i>) symptomatic dogs before and following a 4-week treatment with miltefosine and allopurinol: a double-blin controlled and cross-sectional study. <u>Acta Vet Scand. 65 (1): 2.</u></li> </ol>
Flow Cyte		Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul 1. Barraud-Lange, V. <i>et al.</i> (2020) Partial Sperm beta1 Integrin Subunit Deletion Prove
		Test results have shown that this antibody is also suitable for use as a negative contro with porcine and canine cells.
		recommended for use as a negative control to assess the level of non-specific binding rat IgG2a test antibodies to the surface of human and mouse cells in flow cytometry.

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint