

Datasheet: MCA6035

Description:	MOUSE ANTI MONO- AND POLYUBIQUITIN
Specificity:	MONO- AND POLYUBIQUITIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	FK2
Isotype:	IgG1
Quantity:	50 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
ELISA	▪			
Immunoprecipitation	▪			
Western Blotting	▪			1/100 - 1/1000
Immunocytochemistry	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Broad
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by ion exchange chromatography from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.1% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Poly-ubiquitinated lysozyme

Specificity **Mouse anti mono- and polyubiquitin antibody**, clone FK2 recognizes both mono- and polyubiquitin chains. The antibody does not recognize free ubiquitin.

Ubiquitin, as the name implies, is a ubiquitously expressed and highly conserved protein of 8.6 kDa. The protein is covalently linked to selected lysine residues in a post-translational modification process known as ubiquitylation or ubiquitination. This chemical reaction is mediated by three different protein families; ubiquitin-activating enzymes (also known as E1s), ubiquitin-conjugating enzymes (also known as E2s) and ubiquitin ligases (also known as E3s) ([Hershko and Ciechanover 1998](#)).

The impact of ubiquitination depends on whether a single ubiquitin moiety (monoubiquitination) or an ubiquitin chain (polyubiquitination) has been attached to a protein. Monoubiquitination tends to trigger cellular processes related to endocytosis and membrane trafficking ([Haglund et al. 2003](#)) while the impact of polyubiquitination varies depending on how the ubiquitin residues in the chain have been linked. Attachment of Lysine-48 ubiquitin chains results in degradation by the 20S proteasome while addition of Lysine-63 ubiquitin chains mediates DNA damage and NFkappaB signaling ([Chen 2005](#); [Mocciaro and Rape 2012](#)). Lysine-6, Lysine-11, Lysine-27, Lysine-29 and Lysine-33 chains have also been reported ([Komander 2009](#); [Ye and Rape 2011](#)).

When comparing staining of mouse anti-mono and polyubiquitin antibody (clone FK2) against staining with mouse anti polyubiquitin antibody (clone FK1) one can determine if a protein target is mono- or polyubiquitinated.

Western Blotting Mouse anti mono- and poly-ubiquitin recognizes mono- and poly-ubiquitylated protein conjugates but does not recognize free ubiquitin by Western Blot. Use of milk based blocking reagents is not recommended. 1% BSA in PBS or TBS Tween should be used instead.

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...) [HRP](#)

Rabbit Anti Mouse IgG (STAR12...)	RPE
Rabbit Anti Mouse IgG (STAR8...)	DyLight@800
Goat Anti Mouse IgG (STAR76...)	RPE
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight@488 , DyLight@680 , DyLight@800 , FITC , HRP
Goat Anti Mouse IgG (STAR70...)	FITC

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M382722:210513'

Printed on 18 May 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)