

Datasheet: MCA6037

| Description:         | MOUSE ANTI RelB          |
|----------------------|--------------------------|
| Specificity:         | RelB                     |
| Other names:         | IREL, I-Rel              |
| Format:              | Purified                 |
|                      |                          |
| Product Type:        | Monoclonal Antibody      |
| Product Type: Clone: | Monoclonal Antibody 17.3 |
|                      | ,                        |

# **Product Details**

### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="https://www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

|                            | Yes | No | Not Determined | Suggested Dilution |
|----------------------------|-----|----|----------------|--------------------|
| Immunohistology - Paraffin | -   |    |                | 1/10 - 1/50        |
| Immunoprecipitation        | -   |    |                |                    |
| Western Blotting           | -   |    |                | 1 - 2 ug/ml        |
| Immunofluorescence         | •   |    |                | 1 ug/ml            |
| Immunocytochemistry        | •   |    |                | 1 ug/ml            |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

| Target Species                    | Human   |
|-----------------------------------|---|
| Species Cross<br>Reactivity       | Reacts with: Rhesus Monkey, Mouse  N.B. Antibody reactivity and working conditions may vary between species |
| Product Form                      | Purified IgG - liquid   |
| Preparation                       | Purified IgG prepared by affinity chromatography on Protein A   |
| Buffer Solution                   | Phosphate buffered saline   |
| Preservative<br>Stabilisers       | 0.1% Sodium Azide (NaN <sub>3</sub> )   |
| Approx. Protein<br>Concentrations | IgG concentration 0.5 mg/ml   |
| Immunogen                         | Recombinant protein derived from C-terminal region of human RelB protein                                    |

### External Database Links

UniProt:

Q01201 Related reagents

**Entrez Gene:** 

5971 RELB Related reagents

### Specificity

**Mouse anti-RelB antibody, clone 17.3** recognizes the transcription factor RelB, also known as I-Rel. RelB is encoded by the v-rel avian reticuloendotheliosis viral oncogene homolog B gene and is a critical component of the NF-kappa-B pathway.

Upon cleavage of the NF-kappa-B p105 and p100 subunits into the NF-kappa-B p50 and p52 subunits, RelB forms RelB-p50 and RelB-p52 heterodimers. These dimers act as activators of transcription by binding to kappaB sites present on DNA (<u>Oeckinghaus and Ghosh 2009</u>). RelB-p50 and RelB-p52 binding induces the expression of genes mediating immune responses, inflammation and apoptosis (<u>Jing and Lee 2014</u>).

Misregulation of the NF-kappa-B pathway has been implicated in a variety of disease states including autoimmunity and cancer. As the NF-kappa-B pathway has been demonstrated to be constitutively active in a large group of tumors, developing NF-kappa-B inhibitors has emerged as a new cancer therapy approach (Chaturverdi et al. 2011 and Baud and Karin et al. 2009).

### Western Blotting

Mouse anti RelB antibody recognizes a band of approximately 63 kDa in THP-1 cell lysate, Daudi cell lysate, and Raji cell lysate

### **Storage**

Store at +4°C or at -20°C if preferred.

Storage in frost-free freezers is not recommended.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before

use.

### Guarantee

12 months from date of despatch

# Health And Safety Information

Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf

Regulatory

For research purposes only

## Related Products

## **Recommended Secondary Antibodies**

Goat Anti Mouse IgG IgA IgM (STAR87...) Alk. Phos., HRP

Goat Anti Mouse IgG (STAR77...)

Rabbit Anti Mouse IgG (STAR12...)

RPE

Rabbit Anti Mouse IgG (STAR8...) DyLight®800

Rabbit Anti Mouse IgG (STAR13...)

Goat Anti Mouse IgG (STAR76...)

Goat Anti Mouse IgG (STAR70...)

Goat Anti Mouse IgG (Fc) (STAR120...)

Rabbit Anti Mouse IgG (STAR9...)

FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®680, DyLight®800, FITC, HRP

# **Recommended Negative Controls**

# MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South Tel: +1 800 265 7376

Worldwide

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +1 919 878 3751 America

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_us@bio-rad.com

Email: antibody\_sales\_uk@bio-rad.com

Email: antibody\_sales\_de@bio-rad.com

'M357896:190911'

### Printed on 11 Oct 2019

© 2019 Bio-Rad Laboratories Inc | Legal | Imprint