

Datasheet: MCA6044F

Description:	RAT ANTI MOUSE TER-119/ERYTHROID CELLS:FITC
Specificity:	TER-119/ERYTHROID CELLS
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	TER-119
Isotype:	IgG2b
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/5

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse						
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid						
Max Ex/Em	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>FITC</td> <td>490</td> <td>525</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	FITC	490	525
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
FITC	490	525					
Preparation	Purified IgG prepared by affinity chromatography						
Buffer Solution	Phosphate buffered saline						
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)						
Approx. Protein Concentrations	IgG concentration 0.5 mg/ml						
Immunogen	Day 14 fetal liver cells from a C57BL/6 mouse						

Specificity

Rat anti Mouse TER-119/erythroid cells antibody, clone TER-119 recognizes TER-119 also known as Ly-76. The TER-119 clone detects a 52 kDa protein that is associated with cell surface expressed glycoprotein A. TER-119 is an erythroid-specific antigen expressed from the early proerythroblast through to the mature erythrocyte stage. This antigen is also expressed on 20-25% of bone marrow cells and 2-3% of spleen cells but not on thymocytes, lymph node cells or erythroid

colony-forming cells (Kina *et al.* 2000).

Flow Cytometry	Use 10 µl of the suggested working dilution to label 1x10 ⁶ cells in 100ul
-----------------------	---

Further Reading	1. Kina, T. <i>et al.</i> (2000) The monoclonal antibody TER-119 recognizes a molecule associated with glycophorin A and specifically marks the late stages of murine erythroid lineage. Br J Haematol. 109 (2): 280-7.
------------------------	---

Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.
----------------	--

Guarantee	18 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M344043:190111'

Printed on 20 May 2019

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)