

## Datasheet: MCA6066F

<b>Description:</b>	HAMSTER ANTI MOUSE Fc EPSILON R1 ALPHA:FITC
<b>Specificity:</b>	Fc EPSILON R1 ALPHA
<b>Format:</b>	FITC
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	MAR-1
<b>Isotype:</b>	IgG
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/20 - 1/80

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse						
<b>Product Form</b>	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid						
<b>Max Ex/Em</b>	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>FITC</td> <td>490</td> <td>525</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	FITC	490	525
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
FITC	490	525					
<b>Preparation</b>	Purified IgG prepared by affinity chromatography						
<b>Buffer Solution</b>	Phosphate buffered saline						
<b>Preservative Stabilisers</b>	<0.1% Sodium Azide (NaN <sub>3</sub> )						
<b>Approx. Protein Concentrations</b>	IgG concentration 0.5 mg/ml						
<b>Immunogen</b>	Mouse Fc epsilon receptor alpha subunit						

**External Database  
Links**

**UniProt:**

[P20489](#)   [Related reagents](#)

**Entrez Gene:**

[14125](#)   Fcer1a   [Related reagents](#)

---

**Synonyms**

Fce1a

---

**Specificity**

**Hamster anti mouse Fc epsilon R1 alpha, clone MAR-1** recognizes Fc receptor epsilon R1 alpha also known as FceRIalpha or high affinity IgE receptor. The Fc epsilon R1 complex is involved in initiating IgE-mediated allergic reactions. Fc receptor epsilon is primarily expressed on mast cells and basophils. Activation of Fc epsilon R1 leads to release of allergic mediators such as histamine.

---

**Flow Cytometry**

Use 10 µl of the suggested working dilution to label 1x10<sup>6</sup> cells in 100ul

---

**Storage**

Store at +4°C. DO NOT FREEZE.  
This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee**

12 months from date of despatch

---

**Health And Safety  
Information**

Material Safety Datasheet documentation #10040 available at:  
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

---

**Regulatory**

For research purposes only

---

## Related Products

### Recommended Negative Controls

[HAMSTER \(ARMENIAN\) IgG NEGATIVE CONTROL:FITC \(MCA2356F\)](#)

**North & South**   Tel: +1 800 265 7376

**America**   Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M389695:210806'

Printed on 29 Aug 2021