

Datasheet: MCA6066GA

| | |
|----------------------|--|
| Description: | HAMSTER ANTI MOUSE Fc EPSILON R1 ALPHA |
| Specificity: | Fc EPSILON R1 ALPHA |
| Format: | Purified |
| Product Type: | Monoclonal Antibody |
| Clone: | MAR-1 |
| Isotype: | IgG |
| Quantity: | 0.1 mg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry | ■ | | | 1/20 - 1/80 |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

| | |
|---------------------------------------|---|
| Target Species | Mouse |
| Product Form | Purified IgG - liquid |
| Preparation | Purified IgG prepared by affinity chromatography |
| Buffer Solution | Phosphate buffered saline |
| Preservative Stabilisers | 0.09% Sodium Azide (NaN ₃) |
| Approx. Protein Concentrations | IgG concentration 0.5 mg/ml |
| Immunogen | Mouse Fc epsilon receptor alpha subunit |
| External Database Links | <p>UniProt: P20489 Related reagents</p> <p>Entrez Gene: 14125 Fcer1a Related reagents</p> |
| Synonyms | Fce1a |

| | |
|--------------------------------------|---|
| Specificity | Hamster anti mouse Fc epsilon R1 alpha, clone MAR-1 recognizes Fc receptor epsilon R1 alpha also known as FcεRIalpha or high affinity IgE receptor. The Fc epsilon R1 complex is involved in initiating IgE-mediated allergic reactions. Fc receptor epsilon is primarily expressed on mast cells and basophils. Activation of Fc epsilon R1 leads to release of allergic mediators such as histamine. |
| Flow Cytometry | Use 10 µl of the suggested working dilution to label 1x10 ⁶ cells in 100ul |
| Storage | Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use. |
| Guarantee | 18 months from date of despatch |
| Health And Safety Information | Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf |
| Regulatory | For research purposes only |

Related Products

Recommended Secondary Antibodies

Goat Anti Hamster IgG (STAR104...) [DyLight®800](#), [FITC](#)

Goat Anti Hamster IgG (STAR79...) [Biotin](#), [FITC](#), [HRP](#)

Recommended Negative Controls

[HAMSTER \(ARMENIAN\) IgG NEGATIVE CONTROL \(MCA2356\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M345644:190125'

Printed on 11 Oct 2019

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)