

Datasheet: MCA6076PE

Description:	RAT ANTI MOUSE CD86:RPE
Specificity:	CD86
Other names:	B7-2
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	GL-1
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/2

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse						
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - liquid						
Max Ex/Em	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>RPE 488nm laser</td> <td>496</td> <td>578</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	RPE 488nm laser	496	578
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
RPE 488nm laser	496	578					
Preparation	Purified IgG prepared by affinity chromatography						
Buffer Solution	Phosphate buffered saline						
Preservative Stabilisers	0.09% Sodium Azide						
Approx. Protein Concentrations	IgG concentration 0.2 mg/ml						
Immunogen	LPS-activated CBA/Ca mouse splenic B lymphocytes						
External Database Links	UniProt: P42082 Related reagents						

Entrez Gene:[12524](#) Cd86 [Related reagents](#)

Specificity	Rat anti mouse CD86 antibody, clone GL-1 recognizes mouse CD86 (B7-2), a ~80 kDa cell surface glycoprotein, and a member of the CD28/B7 family. In mouse CD86 is expressed at high levels on peripheral blood monocytes and dendritic cells, and at low levels on resting B and T lymphocytes. Expression of CD86 on these cell populations can be increased upon activation. CD86 has been identified, along with CD80 (B7-1), as a ligand for CD28 and cytotoxic T lymphocyte antigen-4 (CTLA4). CD28 and CTLA4 are two receptors that have opposing functions in T lymphocyte stimulation. Interaction of CD86 with CD28 promotes a number of T lymphocyte activities, whereas the binding of CD86 to CTLA4 inhibits T lymphocyte responses.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul
References	1. Pechhold, K. <i>et al.</i> (1997) Inflammatory cytokines IFN-gamma plus TNF-alpha induce regulated expression of CD80 (B7-1) but not CD86 (B7-2) on murine fibroblasts. J Immunol. 158 (10): 4921-9.
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:RPE \(MCA1212PE\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M344229:190111'

Printed on 11 Oct 2019