Datasheet: MCA6098F

Description:	MOUSE ANTI PIG CD1:FITC
Specificity:	CD1
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	76-7-4
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species

Pig

Product Form

Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
FITC	490	525

Preparation

Purified IgG prepared by combination of precipitation and chromatography techniques

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers

<0.1% Sodium Azide (NaN_3)

Approx. Protein Concentrations

IgG concentration 0.5 mg/ml

Immunogen

Fresh dd miniature swine thymocytes

External Database Links	**UniProt:** Q9XS72 Related reagents **Entrez Gene:** 396785 CD1.1 Related reagents
Synonyms	CD1.1
Specificity	**Mouse anti Pig CD1 antibody, clone 76-7-4** recognizes a porcine homologue of the human CD1 cell surface receptor. CD1 is a member of the immunoglobulin superfamily. While it is structurally similar to the MHC class I receptor CD1 differs functionally by presenting lipid antigens to T cells (Germain & Margulies, 1993). To date, five classes of the CD1 genes have been identified across different mammalian species, namely CD1A, CD1B, CD1C, CD1D, and CD1E (Brigl & Brenner, 2004). In swine various tissues express CD1C even though its gene has been reported to carry a loss-of-function mutation (Eguchi-Ogawa *et al.* 2007). CD1 is predominantly found on dendritic cells, macrophages, monocytes, and thymocytes (Chun *et al.* 1999).
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10^6 cells in 100ul
References	1. Pauly, T. *et al.* (1998) Infection with classical swine fever virus: effects on phenotype and immune responsiveness of porcine T lymphocytes. J Gen Virol. 79 (Pt 1): 31-40.
Storage	Store at +4^{o}C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA6098F 10040
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG2a NEGATIVE CONTROL:FITC (MCA929F)

Recommended Useful Reagents

MOUSE ANTI PIG CD45RA:RPE (MCA1751PE)

North & South America Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M387523:210629'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint