

Datasheet: MCA609AMO

## **BATCH NUMBER 149151**

Description:	RAT ANTI MOUSE CD8 ALPHA:Amethyst Orange
Specificity:	CD8 ALPHA
Other names:	LY-2
Format:	Amethyst Orange
Product Type:	Monoclonal Antibody
Clone:	KT15
Isotype:	lgG2a
Quantity:	0.1 mg

# **Product Details**

## **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Product Form	Purified IgG conjugat	ed to Amethyst Orang	e - liquid
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	Amethyst Orange	405	540
Preparation	Purified IgG prepared supernatant	I by affinity chromatog	raphy on Protein G
uffer Solution	Phosphate buffered s	aline	
Preservative	0.09% Sodium Azide	(NaN <sub>3</sub> )	
Stabilisers	1% Bovine Serum Alk	oumin	
Approx. Protein	IgG concentration 0.1	mg/ml	

Concentrations		
Immunogen	T cell clone, C6	
External Database Links	UniProt: P01731 Related reagents	
	Entrez Gene:  12525 Cd8a Related reagents	
Synonyms	Lyt2, Lyt-2	
Fusion Partners	Spleen cells from immunized SD rats were fused with cells of the NS0 mouse myeloma cell line	
Specificity	Rat anti mouse CD8α, clone KT15, recognizes the <u>alpha chain of mouse CD8</u> . CD8 is a heterodimeric protein composed of disulphide-linked CD8α and <u>CD8β</u> chains that is expressed primarily on cytotoxic T-cells. CD8 functions in the interaction with MHC Class I-bearing targets and plays a role in T-cell-mediated killing ( <u>Nakauchi</u> , H. et al., 1985) & Nakauchi, H. et al., 1987).	

Clone KT15 is reported to block T-cell-mediated cytotoxicity in in vitro assays (Zeis, M. et al., 2002).

#### Flow Cytometry

Use 10ul of the suggested working dilution to label 10<sup>6</sup> cells in 100ul.

The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (BUF041A/B).

#### References

- 1. Tomonari, K. & Lovering, E. (1988) T-cell receptor-specific monoclonal antibodies against a V beta 11-positive mouse T-cell clone. Immunogenetics. 28 (6): 445-51.
- 2. Whiteland, J.L. et al. (1995) Immunohistochemical detection of T-cell subsets and other leukocytes in paraffin-embedded rat and mouse tissues with monoclonal antibodies. J Histochem Cytochem. 43 (3): 313-20.
- 3. Lee, Y.L. et al (2003) Oral administration of Agaricus blazei (H1 strain) inhibited tumor growth in a sarcoma 180 inoculation model. Exp Anim. 52: 371-5.
- 4. Eller, K. et al. (2011) IL-9 production by regulatory T cells recruits mast cells that are essential for regulatory T cell-induced immune suppression. J Immunol. 186: 83-91.
- 5. Grimm, M. et al. (2010) Evaluation of immunological escape mechanisms in a mouse model of colorectal liver metastases. BMC Cancer. 10: 82.
- 6. Liao, D. et al. (2009) Cancer Associated Fibroblasts Promote Tumor Growth and Metastasis by Modulating the Tumor Immune Microenvironment in a 4T1 Murine Breast Cancer Model PLoS One. 4: e7965.
- 7. Moos, M.P. et al. (2005) The lamina adventitia is the major site of immune cell accumulation in standard chow-fed apolipoprotein E-deficient mice. Arterioscler Thromb Vasc Biol. 25: 2386-91.
- 8. Stevenson, P.G. et al. (2002) Uncoupling of virus-induced inflammation and anti-viral

- immunity in the brain parenchyma. J Gen Virol. 83: 1735-43.
- 9. Wang, X. *et al.* (2011) Quercetin and Bornyl Acetate Regulate T-Lymphocyte Subsets and INF-γ/IL-4 Ratio In Utero in Pregnant Mice. <u>Evid Based Complement Alternat Med.</u> 2011: 745262.
- 10. Zeis, M. *et al.* (2002) Idiotype protein-pulsed dendritic cells produce strong anti-myeloma effects after syngeneic stem cell transplantation in mice. <u>Bone Marrow Transplant.</u> 29: 213-21.
- 11. Ideguchi, M. *et al.* (2008) Immune or inflammatory response by the host brain suppresses neuronal differentiation of transplanted ES cell-derived neural precursor cells. J Neurosci Res. 86: 1936-43.
- 12. Wolf, D. *et al.* (2005) CD4+CD25+ regulatory T cells inhibit experimental anti-glomerular basement membrane glomerulonephritis in mice. <u>J Am Soc Nephrol. 16:</u> 1360-70.
- 13. Severinova, J. *et al.* (2005) Co-inoculation of Borrelia afzelii with tick salivary gland extract influences distribution of immunocompetent cells in the skin and lymph nodes of mice. Folia Microbiol (Praha). 50: 457-63.
- 14. Zaini, J. *et al.* (2007) OX40 ligand expressed by DCs costimulates NKT and CD4+ Th cell antitumor immunity in mice. <u>J Clin Invest. 117: 3330-8.</u>
- 15. Meyer, C. *et al.* (2011) Chronic inflammation promotes myeloid-derived suppressor cell activation blocking antitumor immunity in transgenic mouse melanoma model. <u>Proc Natl</u> Acad Sci U S A. 108: 17111-6.
- 16. Zitt, E. *et al.* (2011) The selective mineralocorticoid receptor antagonist eplerenone is protective in mild anti-GBM glomeru-lonephritis. Int J Clin Exp Pathol. 4:606-15.
- 17. Singh, V. *et al.* (2011) Co-administration of IL-1+IL-6+TNF-α with Mycobacterium tuberculosis infected macrophages vaccine induces better protective T cell memory than BCG. PLoS One. 6: e16097.
- 18. Kalyanasundaram Bhanumathy, K. *et al.* (2015) Potent immunotherapy against well-established thymoma using adoptively transferred transgene IL-6-engineered dendritic cell-stimulated CD8(+) T-cells with prolonged survival and enhanced cytotoxicity. <u>J Gene Med. 17 (8-9): 153-60.</u>
- 19. Abiko K *et al.* (2015) IFN-γ from lymphocytes induces PD-L1 expression and promotes progression of ovarian cancer. <u>Br J Cancer.</u> 112 (9): 1501-9.
- 20. Phan-Lai, V. *et al.* (2016) The Antitumor Efficacy of IL2/IL21-Cultured Polyfunctional Neu-Specific T Cells Is TNFα/IL17 Dependent. <u>Clin Cancer Res. 22 (9): 2207-16.</u>
- 21. Kajiwara, T. *et al.* (2016) Hypoxia augments MHC class I antigen presentation via facilitation of ERO1-α-mediated oxidative folding in murine tumor cells. <u>Eur J Immunol.</u> <u>Sep 26. [Epub ahead of print]</u>
- 22. Srivastava, A.K. *et al.* (2016) Co-transplantation of syngeneic mesenchymal stem cells improves survival of allogeneic glial-restricted precursors in mouse brain. <u>Exp Neurol. 275 Pt 1: 154-61.</u>
- 23. Meier, R.P. *et al.* (2014) Survival of free and encapsulated human and rat islet xenografts transplanted into the mouse bone marrow. <u>PLoS One. 9 (3): e91268.</u>
- 24. Groh, J. *et al.* (2021) Immune modulation attenuates infantile neuronal ceroid lipofuscinosis in mice before and after disease onset <u>Brain Communications</u>. fcab047 [Epub ahead of print].

**Storage** 

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA609AMO">https://www.bio-rad-antibodies.com/SDS/MCA609AMO</a>
Regulatory	For research purposes only

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M375939:210114'

### Printed on 15 Aug 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint