

Datasheet: MCA609SBUV795

BATCH NUMBER 100006151

Description:	RAT ANTI MOUSE CD8 ALPHA:StarBright UltraViolet 795
Specificity:	CD8 ALPHA
Other names:	LY-2
Format:	StarBright UltraViolet 795
Product Type:	Monoclonal Antibody
Clone:	KT15
Isotype:	IgG2a
Quantity:	100 TESTS/0.5ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse						
Product Form	Purified IgG conjugated to StarBright UltraViolet 795 - liquid						
Max Ex/Em	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>StarBright UltraViolet 795</td> <td>340</td> <td>792</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	StarBright UltraViolet 795	340	792
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
StarBright UltraViolet 795	340	792					
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant						
Buffer Solution	Phosphate buffered saline						
Preservative	0.09% Sodium Azide (NaN ₃)						
Stabilisers	1% Bovine Serum Albumin 0.1% Pluronic F68						

0.1% PEG 3350
0.05% Tween 20

Immunogen T cell clone, C6

External Database Links

UniProt:

[P01731](#) [Related reagents](#)

Entrez Gene:

[12525](#) Cd8a [Related reagents](#)

Synonyms Lyt2, Lyt-2

Fusion Partners Spleen cells from immunized SD rats were fused with cells of the NS0 mouse myeloma cell line

Specificity **Rat anti Mouse CD8 α , clone KT15**, recognizes the [alpha chain of mouse CD8](#). CD8 is a heterodimeric protein composed of disulphide-linked CD8 α and [CD8 \$\beta\$](#) chains that is expressed primarily on cytotoxic T-cells. CD8 functions in the interaction with MHC Class I-bearing targets and plays a role in T-cell-mediated killing ([Nakauchi, H. et al., 1985](#) & [Nakauchi, H. et al., 1987](#)).

Clone KT15 is reported to block T-cell-mediated cytotoxicity in *in vitro* assays ([Zeis, M. et al., 2002](#)).

Flow Cytometry Use 5ul of the suggested working dilution to label 10⁶ cells in 100ul. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.

References

1. Tomonari, K. & Lovering, E. (1988) T-cell receptor-specific monoclonal antibodies against a V beta 11-positive mouse T-cell clone. [Immunogenetics. 28 \(6\): 445-51.](#)
2. Whiteland, J.L. et al. (1995) Immunohistochemical detection of T-cell subsets and other leukocytes in paraffin-embedded rat and mouse tissues with monoclonal antibodies. [J Histochem Cytochem. 43 \(3\): 313-20.](#)
3. Lee, Y.L. et al (2003) Oral administration of Agaricus blazei (H1 strain) inhibited tumor growth in a sarcoma 180 inoculation model. [Exp Anim. 52: 371-5.](#)
4. Eller, K. et al. (2011) IL-9 production by regulatory T cells recruits mast cells that are essential for regulatory T cell-induced immune suppression. [J Immunol. 186: 83-91.](#)
5. Grimm, M. et al. (2010) Evaluation of immunological escape mechanisms in a mouse model of colorectal liver metastases. [BMC Cancer. 10: 82.](#)
6. Liao, D. et al. (2009) Cancer Associated Fibroblasts Promote Tumor Growth and Metastasis by Modulating the Tumor Immune Microenvironment in a 4T1 Murine Breast Cancer Model [PLoS One. 4: e7965.](#)
7. Moos, M.P. et al. (2005) The lamina adventitia is the major site of immune cell accumulation in standard chow-fed apolipoprotein E-deficient mice. [Arterioscler Thromb Vasc Biol. 25: 2386-91.](#)
8. Stevenson, P.G. et al. (2002) Uncoupling of virus-induced inflammation and anti-viral immunity in the brain parenchyma. [J Gen Virol. 83: 1735-43.](#)

9. Wang, X. *et al.* (2011) Quercetin and Bornyl Acetate Regulate T-Lymphocyte Subsets and INF- γ /IL-4 Ratio In Utero in Pregnant Mice. [Evid Based Complement Alternat Med. 2011: 745262.](#)
10. Zeis, M. *et al.* (2002) Idiotype protein-pulsed dendritic cells produce strong anti-myeloma effects after syngeneic stem cell transplantation in mice. [Bone Marrow Transplant. 29: 213-21.](#)
11. Ideguchi, M. *et al.* (2008) Immune or inflammatory response by the host brain suppresses neuronal differentiation of transplanted ES cell-derived neural precursor cells. [J Neurosci Res. 86: 1936-43.](#)
12. Wolf, D. *et al.* (2005) CD4+CD25+ regulatory T cells inhibit experimental anti-glomerular basement membrane glomerulonephritis in mice. [J Am Soc Nephrol. 16: 1360-70.](#)
13. Severinova, J. *et al.* (2005) Co-inoculation of *Borrelia afzelii* with tick salivary gland extract influences distribution of immunocompetent cells in the skin and lymph nodes of mice. [Folia Microbiol \(Praha\). 50: 457-63.](#)
14. Zaini, J. *et al.* (2007) OX40 ligand expressed by DCs costimulates NKT and CD4+ Th cell antitumor immunity in mice. [J Clin Invest. 117: 3330-8.](#)
15. Meyer, C. *et al.* (2011) Chronic inflammation promotes myeloid-derived suppressor cell activation blocking antitumor immunity in transgenic mouse melanoma model. [Proc Natl Acad Sci U S A. 108: 17111-6.](#)
16. Zitt, E. *et al.* (2011) The selective mineralocorticoid receptor antagonist eplerenone is protective in mild anti-GBM glomerulonephritis. [Int J Clin Exp Pathol. 4:606-15.](#)
17. Singh, V. *et al.* (2011) Co-administration of IL-1+IL-6+TNF- α with Mycobacterium tuberculosis infected macrophages vaccine induces better protective T cell memory than BCG. [PLoS One. 6: e16097.](#)
18. Kalyanasundaram Bhanumathy, K. *et al.* (2015) Potent immunotherapy against well-established thymoma using adoptively transferred transgene IL-6-engineered dendritic cell-stimulated CD8(+) T-cells with prolonged survival and enhanced cytotoxicity. [J Gene Med. 17 \(8-9\): 153-60.](#)
19. Abiko K *et al.* (2015) IFN- γ from lymphocytes induces PD-L1 expression and promotes progression of ovarian cancer. [Br J Cancer. 112 \(9\): 1501-9.](#)
20. Phan-Lai, V. *et al.* (2016) The Antitumor Efficacy of IL2/IL21-Cultured Polyfunctional Neu-Specific T Cells Is TNF α /IL17 Dependent. [Clin Cancer Res. 22 \(9\): 2207-16.](#)
21. Kajiwara, T. *et al.* (2016) Hypoxia augments MHC class I antigen presentation via facilitation of ERO1- α -mediated oxidative folding in murine tumor cells. [Eur J Immunol. 46 \(12\): 2842-51.](#)
22. Srivastava, A.K. *et al.* (2016) Co-transplantation of syngeneic mesenchymal stem cells improves survival of allogeneic glial-restricted precursors in mouse brain. [Exp Neurol. 275 Pt 1: 154-61.](#)
23. Meier, R.P. *et al.* (2014) Survival of free and encapsulated human and rat islet xenografts transplanted into the mouse bone marrow. [PLoS One. 9 \(3\): e91268.](#)
24. Groh, J. *et al.* (2021) Immune modulation attenuates infantile neuronal ceroid lipofuscinosis in mice before and after disease onset [Brain Communications. 3\(2\): fcab047.](#)
25. Cecil, D.L. *et al.* (2022) COX-2 inhibitors decrease expression of PD-L1 in colon tumors and increase the influx of Type I tumor infiltrating lymphocytes. [Cancer Prev Res \(Phila\). canprevres.0227.2021.](#)

26. Karikari, A.A. *et al.* (2022) Neurodegeneration by α -synuclein-specific T cells in AAV-A53T- α -synuclein Parkinson's disease mice. [Brain Behav Immun. 101: 194-210.](#)
27. Badr, M. *et al.* (2022) Expansion of regulatory T cells by CD28 superagonistic antibodies attenuates neurodegeneration in A53T- α -synuclein Parkinson's disease mice. [J Neuroinflammation. 19 \(1\): 319.](#)
28. McFleder, R.L. *et al.* (2023) Brain-to-gut trafficking of alpha-synuclein by CD11c(+) cells in a mouse model of Parkinson's disease. [Nat Commun. 14 \(1\): 7529.](#)
29. Aloui, A. *et al.* (2023) AFM₁ Exposure in Male Balb/c Mice and Intervention Strategies Against Its Immuno-physiological toxicity using Clay Mineral and Lactic Acid Bacteria Alone or in Combination. [Immunopharmacol Immunotoxicol. : 1-32.](#)
30. Aringer, I. *et al.* (2021) Agonism of Prostaglandin E2 Receptor 4 Ameliorates Tubulointerstitial Injury in Nephrotoxic Serum Nephritis in Mice. [J Clin Med. 10 \(4\) :832.](#)

Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.
Guarantee	12 months from date of despatch
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts
Health And Safety Information	Material Safety Datasheet documentation #20471 available at: https://www.bio-rad-antibodies.com/SDS/MCA609SBUV79520471
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M399315:220630'

Printed on 08 Mar 2024