

Datasheet: MCA609SBY665 BATCH NUMBER 100007711

Description:	RAT ANTI MOUSE CD8 ALPHA:StarBright Yellow 665			
Specificity:	CD8 ALPHA			
Other names:	LY-2			
Format:	StarBright Yellow 665			
Product Type:	Monoclonal Antibody			
Clone:	KT15			
Isotype:	lgG2a			
Quantity:	100 TESTS/0.5ml			

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
		Yes No	Not Determined	Suggested Dilution		
	Flow Cytometry	•		Neat		
	necessarily exclude its	use in such proced nmended that the us	use in a particular tech ures. Suggested working ser titrates the product fo controls.	g dilutions are given as		
Target Species	Mouse					
Product Form	Purified IgG conjugate	d to StarBright Yello	w 665 - liquid			
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)			
	StarBright Yellow 665	554	670			
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant					
Buffer Solution	Phosphate buffered saline					
Preservative Stabilisers	0.09% Sodium Azide (l 1% Bovine Serum Albu 0.1% Pluronic F68 0.1% PEG 3350	0,				

	0.05% Tween 20
Immunogen	T cell clone, C6
External Database Links	UniProt: <u>P01731</u> <u>Related reagents</u> Entrez Gene: <u>12525</u> Cd8a <u>Related reagents</u>
Synonyms	Lyt2, Lyt-2
Fusion Partners	Spleen cells from immunized SD rats were fused with cells of the NS0 mouse myeloma cell line
Specificity	Rat anti Mouse CD8α, clone KT15, recognizes the <u>alpha chain of mouse CD8</u> . CD8 is a heterodimeric protein composed of disulphide-linked CD8α and <u>CD8β</u> chains that is expressed primarily on cytotoxic T-cells. CD8 functions in the interaction with MHC Class I-bearing targets and plays a role in T-cell-mediated killing (<u>Nakauchi , H. <i>et al.</i>, 1985</u> & <u>Nakauchi, H. <i>et al.</i>, 1987</u>).
	Clone KT15 is reported to block T-cell-mediated cytotoxicity in <i>in vitro</i> assays (Zeis, M. et <u>al., 2002</u>).
Flow Cytometry	Use 5µl of the suggested working dilution to label 10 ⁶ cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.
References	 Tomonari, K. & Lovering, E. (1988) T-cell receptor-specific monoclonal antibodies against a V beta 11-positive mouse T-cell clone. <u>Immunogenetics. 28 (6): 445-51.</u> Whiteland, J.L. <i>et al.</i> (1995) Immunohistochemical detection of T-cell subsets and other leukocytes in paraffin-embedded rat and mouse tissues with monoclonal antibodies. <u>J</u> <u>Histochem Cytochem. 43 (3): 313-20.</u> Lee, Y.L. <i>et al</i> (2003) Oral administration of Agaricus blazei (H1 strain) inhibited tumor growth in a sarcoma 180 inoculation model. <u>Exp Anim. 52: 371-5.</u> Eller, K. <i>et al.</i> (2011) IL-9 production by regulatory T cells recruits mast cells that are essential for regulatory T cell-induced immune suppression. <u>J Immunol. 186: 83-91.</u> Grimm, M. <i>et al.</i> (2010) Evaluation of immunological escape mechanisms in a mouse model of colorectal liver metastases. <u>BMC Cancer. 10: 82.</u> Liao, D. <i>et al.</i> (2009) Cancer Associated Fibroblasts Promote Tumor Growth and Metastasis by Modulating the Tumor Immune Microenvironment in a 4T1 Murine Breast Cancer Model PLoS One. 4: e7965.

and INF-γ/IL-4 Ratio In Utero in Pregnant Mice. <u>Evid Based Complement Alternat Med.</u> 2011: 745262.

10. Zeis, M. *et al.* (2002) Idiotype protein-pulsed dendritic cells produce strong anti-myeloma effects after syngeneic stem cell transplantation in mice. <u>Bone Marrow</u> <u>Transplant. 29: 213-21.</u>

11. Ideguchi, M. *et al.* (2008) Immune or inflammatory response by the host brain suppresses neuronal differentiation of transplanted ES cell-derived neural precursor cells. <u>J Neurosci Res. 86: 1936-43.</u>

12. Wolf, D. *et al.* (2005) CD4+CD25+ regulatory T cells inhibit experimental anti-glomerular basement membrane glomerulonephritis in mice. <u>J Am Soc Nephrol. 16:</u> <u>1360-70.</u>

13. Severinova, J. *et al.* (2005) Co-inoculation of *Borrelia afzelii* with tick salivary gland extract influences distribution of immunocompetent cells in the skin and lymph nodes of mice. <u>Folia Microbiol (Praha). 50: 457-63.</u>

14. Zaini, J. *et al.* (2007) OX40 ligand expressed by DCs costimulates NKT and CD4+ Th cell antitumor immunity in mice. <u>J Clin Invest. 117: 3330-8.</u>

15. Meyer, C. *et al.* (2011) Chronic inflammation promotes myeloid-derived suppressor cell activation blocking antitumor immunity in transgenic mouse melanoma model. <u>Proc Natl Acad Sci U S A. 108: 17111-6.</u>

16. Zitt, E. *et al.* (2011) The selective mineralocorticoid receptor antagonist eplerenone is protective in mild anti-GBM glomeru-lonephritis. <u>Int J Clin Exp Pathol. 4:606-15.</u>

17. Singh, V. *et al.* (2011) Co-administration of IL-1+IL-6+TNF-α with Mycobacterium tuberculosis infected macrophages vaccine induces better protective T cell memory than BCG. <u>PLoS One. 6: e16097.</u>

18. Kalyanasundaram Bhanumathy, K. *et al.* (2015) Potent immunotherapy against well-established thymoma using adoptively transferred transgene IL-6-engineered dendritic cell-stimulated CD8(+) T-cells with prolonged survival and enhanced cytotoxicity. J Gene Med. 17 (8-9): 153-60.

19. Abiko K *et al.* (2015) IFN-γ from lymphocytes induces PD-L1 expression and promotes progression of ovarian cancer. <u>Br J Cancer. 112 (9): 1501-9.</u>

20. Phan-Lai, V. *et al.* (2016) The Antitumor Efficacy of IL2/IL21-Cultured Polyfunctional Neu-Specific T Cells Is TNFα/IL17 Dependent. <u>Clin Cancer Res. 22 (9): 2207-16.</u>

21. Kajiwara, T. *et al.* (2016) Hypoxia augments MHC class I antigen presentation via facilitation of ERO1- α -mediated oxidative folding in murine tumor cells. <u>Eur J Immunol. 46</u> (12): 2842-51.

22. Srivastava, A.K. *et al.* (2016) Co-transplantation of syngeneic mesenchymal stem cells improves survival of allogeneic glial-restricted precursors in mouse brain. <u>Exp Neurol. 275</u> Pt 1: 154-61.

23. Meier, R.P. *et al.* (2014) Survival of free and encapsulated human and rat islet xenografts transplanted into the mouse bone marrow. <u>PLoS One. 9 (3): e91268.</u>

24. Groh, J. *et al.* (2021) Immune modulation attenuates infantile neuronal ceroid lipofuscinosis in mice before and after disease onset <u>Brain Communications. 3(2):</u> <u>fcab047.</u>

25. Cecil, D.L. *et al.* (2022) COX-2 inhibitors decrease expression of PD-L1 in colon tumors and increase the influx of Type I tumor infiltrating lymphocytes. <u>Cancer Prev Res</u> (<u>Phila</u>). <u>canprevres.0227.2021</u>.

26. Karikari, A.A. et al. (2022) Neurodegeneration by α-synuclein-specific T cells in

	 AAV-A53T-α-synuclein Parkinson's disease mice. Brain Behav Immun. 101: 194-210 27. Badr, M. <i>et al.</i> (2022) Expansion of regulatory T cells by CD28 superagonistic antibodies attenuates neurodegeneration in A53T-α-synuclein Parkinson's disease r Neuroinflammation. 19 (1): 319. 28. McFleder, R.L. <i>et al.</i> (2023) Brain-to-gut trafficking of alpha-synuclein by CD11c cells in a mouse model of Parkinson's disease. Nat Commun. 14 (1): 7529. 29. Aloui, A. <i>et al.</i> (2023) AFM₁ Exposure in Male Balb/c Mice and Intervention Strates Against Its Immuno-physiological toxicity using Clay Mineral and Lactic Acid Bacteria Alone or in Combination. Immunopharmacol Immunotoxicol. : 1-32. 			
	30. Aringer, I. <i>et al.</i> (2021) Agonism of Prostaglandin E2 Receptor 4 Ameliorates Tubulointerstitial Injury in Nephrotoxic Serum Nephritis in Mice. <u>J Clin Med. 10 (4) :832.</u>			
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.			
Guarantee	12 months from date of despatch			
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts			
Health And Safety Information	Material Safety Datasheet documentation #20471 available at: https://www.bio-rad-antibodies.com/SDS/MCA609SBY665 20471			
Regulatory	For research purposes only			

Related Products

Recommended Useful Reagents

MOUSE SEROBLOCK FcR (BUF041A) MOUSE SEROBLOCK FcR (BUF041B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-	rad.com	Email: antibody_sales_uk@bio-	rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M419127:230522'

Printed on 10 May 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint