

Datasheet: MCA6127PE

BATCH NUMBER 164228

Description:	MOUSE ANTI HUMAN CD195:RPE		
Specificity:	CD195		
Other names:	CCR5		
Format:	RPE		
Product Type:	Monoclonal Antibody		
Clone:	T21/8		
Isotype:	lgG1		

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human				
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - liquid				
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm		
	RPE 488nm laser	496	578		
	RPE 561nm laser	546	578		
Preparation	Purified IgG prepared supernatant	by affinity chromatogi	raphy on Protein A		
Buffer Solution	Phosphate buffered sa	aline			
Preservative	<0.1% Sodium Azide	(NaN ₃)			
Stabilisers	0.2% Bovine Serum A	Albumin			

Immunogen	CCR5 peptide (Met1-Lys22) KLH conjugate
External Database Links	UniProt: P51681 Related reagents Entrez Gene: 1234 CCR5 Related reagents
Synonyms	CMKBR5
Specificity	Mouse anti Human CD195, clone T21/8 recognizes CD195 also known as C-C chemokine receptor 5 (CCR5) which belongs to the GPCR protein family and plays a role in inflammation by targeting cells to sites of inflammation. CD195 is expressed in a range of hematopoietic cells including macrophages, dendritic cells and memory T cells. It is also expressed by non-hematopoietic cells including vascular smooth muscle cells and astrocytes (Barmania &Pepper 2013).
	During inflammation CD195 expression is upregulated in CD8+ cells, allowing the cells to migrate to sites with CD4+ T cell and dendritic cell interactions (<u>Castellino et al. 2006</u>). An additional well characterized feature of CD195 is its role as an essential co-receptor for macrophage tropic HIV strains, allowing the virus to enter the target cell during initial infection (<u>Dragic et al. 1996</u>).
	This clone T28/1 has been used in flow cytometry experiments to examine expression levels in CD195 overexpressing PM1 cells. These cells were then used to examine the effect of CD195 expression levels on HIV infections (Monde et al. 2007).
Purity	>95% by SDS PAGE
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light.
Guarantee	Guaranteed for 12 months from the date of despatch or until the date of expiry, whichever comes first. Please see label for expiry date.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA6127PE 10041
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:RPE (MCA928PE)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M401896:220718'

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint