

Datasheet: MCA6127PE

Description:	MOUSE ANTI HUMAN CD195:RPE
Specificity:	CD195
Other names:	CCR5
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	T21/8
Isotype:	IgG1
Quantity:	100 TESTS/1ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
	RPE 561nm laser	546	578
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	<0.1% Sodium Azide (NaN ₃) 0.2% Bovine Serum Albumin		
Immunogen	CCR5 peptide (Met1-Lys22) KLH conjugate		

**External Database
Links**

UniProt:

[P51681](#)

[Related reagents](#)

Entrez Gene:

[1234](#)

CCR5

[Related reagents](#)

Synonyms

CMKBR5

Specificity

Mouse anti Human CD195, clone T21/8 recognizes CD195 also known as C-C chemokine receptor 5 (CCR5) which belongs to the GPCR protein family and plays a role in inflammation by targeting cells to sites of inflammation.

CD195 is expressed in a range of hematopoietic cells including macrophages, dendritic cells and memory T cells. It is also expressed by non-hematopoietic cells including vascular smooth muscle cells and astrocytes ([Barmania & Pepper 2013](#)).

During inflammation CD195 expression is upregulated in CD8+ cells, allowing the cells to migrate to sites with CD4+ T cell and dendritic cell interactions ([Castellino et al. 2006](#)). An additional well characterized feature of CD195 is its role as an essential co-receptor for macrophage tropic HIV strains, allowing the virus to enter the target cell during initial infection ([Dragic et al. 1996](#)).

This clone T28/1 has been used in flow cytometry experiments to examine expression levels in CD195 overexpressing PM1 cells. These cells were then used to examine the effect of CD195 expression levels on HIV infections ([Monde et al. 2007](#)).

Purity

>95% by SDS PAGE

Flow Cytometry

Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul

Storage

This product is shipped at ambient temperature.
Store at +4°C. DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light.

Guarantee

Guaranteed for 12 months from the date of despatch or until the date of expiry, whichever comes first. Please see label for expiry date.

**Health And Safety
Information**

Material Safety Datasheet documentation #10041 available at:
<https://www.bio-rad-antibodies.com/SDS/MCA6127PE>
10041

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:RPE \(MCA928PE\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M441164:250523'

Printed on 23 May 2025

© 2025 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)