

Datasheet: MCA6132F

Description:	MOUSE ANTI HUMAN CD274:FITC		
Specificity:	CD274		
Other names:	PD-L1		
Format:	FITC		
Product Type:	Monoclonal Antibody		
Clone:	29E.2A3		
Isotype:	lgG2b		
Quantity:	100 TESTS/400μl		

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human			
Species Cross	Reacts with: Prima	ate		
Reactivity	N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications personal communications from the originators. Please refer to references indicated further information.			
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid			
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm))
	FITC	490	525	
Preparation	Purified IgG prepa	ared by affinity chromatog	raphy on Protein A f	rom tissue culture
Buffer Solution	Phosphate buffere			

Preservative Stabilisers	<0.1% Sodium Azide (NaN ₃) 0.2% Bovine Serum Albumin
Immunogen	Full length human CD274
External Database Links	UniProt: Q9NZQ7 Related reagents Entrez Gene: 29126 CD274 Related reagents
Synonyms	B7H1, PDCD1L1, PDCD1LG1, PDL1
Specificity	Mouse anti Human CD274 antibody, clone 29E.2A3 detects human CD274, also known as B7-H1 and PD-1L, a cell surface glycoprotein which is a member of the B7 family of co-stimulatory molecules. CD274 is expressed constitutively on macrophages and dendritic cells, and is induced on activated T cells, B cells, endothelial cells and epithelial cells in response to interferons alpha, beta and gamma.
	CD274 is reported to possess dual functions; inhibition of activated effector T cells and co-stimulation of naïve T cells. CD274 inhibits proliferation of activated T cells via ligation to the co-inhibitory molecule CD279 (programmed death-1; PD-1) leading to the secretion of the regulatory cytokine interleukin-10. CD274 has also been shown to co-stimulate early T cell priming and differentiation.
	Deregulated CD274 function has been reported in chronic viral and intracellular bacterial infection, as well as in many autoimmune diseases and cancers.
Purity	>95% by SDS PAGE
Flow Cytometry	Use 4ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light.
Guarantee	Guaranteed for 12 months from the date of despatch or until the date of expiry, whichever comes first. Please see label for expiry date.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA6132F 10041
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG2b NEGATIVE CONTROL:FITC (MCA691F)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M401930:220718'

Printed on 24 Feb 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint