

Datasheet: MCA6133F

BATCH NUMBER 166096

Description:	MOUSE ANTI HUMAN CD279:FITC
Specificity:	CD279
Other names:	PD-1
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	EH12.2H7
Isotype:	IgG1
Quantity:	100 TESTS/400µl

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species

Human

Species Cross Reactivity

Reacts with: Primate

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
FITC	490	525

Preparation

Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

Buffer Solution	Phosphate buffered saline
Preservative	<0.1% Sodium Azide (NaN ₃)
Stabilisers	0.2% Bovine Serum Albumin
External Database Links	<p>UniProt: Q15116 Related reagents</p> <p>Entrez Gene: 5133 PDCD1 Related reagents</p>
Synonyms	PD1
Specificity	<p>Mouse anti Human CD279 antibody, clone EH12.2H7 detects CD279, a co-stimulatory molecule also known as programmed cell death-1 (PD-1).</p> <p>CD279 is a ~50-55 kDa membrane protein which is a member of the CD28 family, and functions as a regulator of T cell activation (Fife and Pauken, 2011). It has two specific ligands; CD274 (PD-L1) and CD273 (PD-L2), and their interaction is key in the balance between stimulatory and inhibitory signals needed for effective immune responses to microbes and self-tolerance.</p> <p>CD279 is a marker for germinal center-associated T cells (Dorfman et al, 2006) and is inducibly expressed by T cells, B cells, NKT cells and monocytes upon activation.</p>
Purity	>95% by SDS PAGE
Flow Cytometry	Use 4ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul
Storage	<p>Store at +4°C. DO NOT FREEZE.</p> <p>This product should be stored undiluted. This product is photosensitive and should be protected from light.</p>
Guarantee	Guaranteed for 12 months from the date of despatch or until the date of expiry, whichever comes first. Please see label for expiry date.
Health And Safety Information	<p>Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA6133F 10041</p>
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

To
find a

batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M401906:220718'

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)