

Datasheet: MCA6133F BATCH NUMBER 170757

Description:	MOUSE ANTI HUMAN CD279:FITC
Specificity:	CD279
Other names:	PD-1
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	EH12.2H7
Isotype:	lgG1
Quantity:	100 TESTS/400µl

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human				
Species Cross Reactivity	•	activity and working conditi	, ,	•	
	•	ed from testing within our land dications from the originated n.		•	
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid				
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm))	
	FITC	490	525	_	
Preparation	Purified IgG prepa	ared by affinity chromatog	raphy on Protein A f	rom tissue culture	

Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	<0.1% Sodium Azide (NaN ₃) 0.2% Bovine Serum Albumin		
External Database Links	UniProt: Q15116 Related reagents		
	Entrez Gene: <u>5133</u> PDCD1 <u>Related reagents</u>		
Synonyms	PD1		
Specificity	Mouse anti Human CD279 antibody, clone EH12.2H7 detects <u>CD279</u> , a co-stimulatory molecule also known as programmed cell death-1 (PD-1).		
	CD279 is a ~50-55 kDa membrane protein which is a member of the CD28 family, and functions as a regulator of T cell activation (<u>Fife and Pauken, 2011</u>). It has two specific ligands; CD274 (PD-L1) and CD273 (PD-L2), and their interaction is key in the balance between stimulatory and inhibitory signals needed for effective immune responses to microbes and self-tolerance.		
	CD279 is a marker for germinal center-associated T cells (<u>Dorfman et al, 2006</u>) and is inducibly expressed by T cells, B cells, NKT cells and monocytes upon activation.		
Purity	>95% by SDS PAGE		
Flow Cytometry	Use 4ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul		
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light.		
Guarantee	Guaranteed for 12 months from the date of despatch or until the date of expiry, whichever comes first. Please see label for expiry date.		
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA6133F		
Regulatory	For research purposes only		

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:FITC (MCA928F)

Product inquiries: www.bio-rad-antibodies.com/technical-support

Printed on 15 Aug 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint