

Datasheet: MCA6135APC

#### **BATCH NUMBER 149889**

Description:	MOUSE ANTI HUMAN CD324:APC
Specificity:	CD324
Other names:	E-Cadherin
Format:	APC
Product Type:	Monoclonal Antibody
Clone:	67A4
Isotype:	lgG1
Quantity:	100 TESTS/1ml

## **Product Details**

### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugate	PC) - liquid	
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	APC	650	661
	<b>.</b>	by anning chromatogi	raphy on Protein A from tissue cult
Buffer Solution	supernatant  Phosphate buffered sa		apny on Protein A from tissue cuit
Buffer Solution Preservative	supernatant	aline	apny on Protein A from tissue cuit
	supernatant  Phosphate buffered sa	aline (NaN <sub>3</sub> )	apny on Protein A from tissue cuit

External Database Links	UniProt:
	P12830 Related reagents
	Entrez Gene:
	999 CDH1 Related reagents
Synonyms	CDHE, UVO
Specificity	<b>Mouse anti Human CD324, clone 67A4</b> recognizes CD324 also known as E-Cadherin, a transmembrane glycoprotein which mediates calcium dependant cell adhesion and cell junction formation.
	The extracellular domain mediates a Ca2+ dependant homophillic interaction with a cadherin molecule on the surface of a neighboring cell. This leads to the formation of tight junctions which assist in blocking the movement of cells and facilitating cell-cell interactions (Gloushankova et al. 2017).
	The cytoplasmic domain is able to interact with catenins which link the CD324 to the actin cytoskeleton which help mediate downstream signaling events growth inhibitory signals to prevent uncontrolled tissue growth during embryonic development, tissue regeneration, wound healing. Loss of CD324 expression has been associated with tumor progression and metastasis. It is thought that metastasis is more likely to occur due to the loss of a maintained stable physical link between tumor cells (Mendonsa et al. 2018).
	This clone 67A4 has been used in immunocytochemistry experiments to examine the localization of CD324 on medullary epithelial cells (Kutleša et al. 2002).
Purity	>95% by SDS PAGE
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul
Storage	Store at +4°C. DO NOT FREEZE.  This product should be stored undiluted. This product is photosensitive and should be protected from light.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA6135APC">https://www.bio-rad-antibodies.com/SDS/MCA6135APC</a> 10041
Regulatory	For research purposes only

# **Related Products**

**Recommended Negative Controls** 

MOUSE IgG1 NEGATIVE CONTROL:APC (MCA928APC)

North & South Tel: +1 800 265 7376 Worldwide Tel: +44 (0)1865 852 700 Europe Tel: +49 (0) 89 8090 95 21 America

Fax: +1 919 878 3751 Fax: +44 (0)1865 852 739 Fax: +49 (0) 89 8090 95 50 Email: antibody\_sales\_de@bio-rad.comd a Email: antibody\_sales\_us@bio-rad.com Email: antibody\_sales\_uk@bio-rad.com

То

batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M353626:190509'

#### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint