

Datasheet: MCA6146F

BATCH NUMBER 161978

Description:	MOUSE ANTI HUMAN CD3:FITC
Specificity:	CD3
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	OKT3
Isotype:	IgG2a
Quantity:	100 TESTS/0.5ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/33

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)		
Approx. Protein Concentrations	IgG concentration 0.2 mg/ml		

Immunogen	Human peripheral blood lymphocytes
External Database Links	<p>UniProt: P07766 Related reagents</p> <p>Entrez Gene: 916 CD3E Related reagents</p>
Synonyms	T3E
Specificity	<p>Mouse anti Human CD3 antibody, clone OKT3, recognizes T-cell surface glycoprotein CD3 epsilon chain (CD3ε), also known as T-cell surface antigen T3/Leu-4 epsilon. CD3ε is a subunit within the multiprotein cell surface receptor TCR-CD3 complex, which is expressed on the surface of mature T cells, NKT cells and thymocytes (Mariuzza et al. 2020). The CD3 complex is required for intracellular signaling and for TCR surface expression. Recruitment of the TCR/CD3 complex initiates early events leading to proliferation and differentiation of mature T cells into an effector cell (Guy & Vignali 2009).</p> <p>Mice that lack the CD3ε gene exhibit early arrest in T cell development and mutation of the gene in humans has been seen to result in severe combined immunodeficiency (SCID) (DeJarnette et al. 1998, de Saint Basile et al. 2004).</p> <p>Mouse anti Human CD3, clone OKT3 is commonly used in flow cytometry in the phenotyping human T cells. Characterization of the OKT3 clone has shown that it is able to compete and block the binding of the CD3 clones SK7 and UCHT1 to human T cells (Li et al. 2005). The OKT3 clone has also been found to have immunosuppressive properties in vivo. As result of this, the clone was humanized to be used as a therapeutic drug called Muromonomab (Orthoclone). Muromonomab was used as an immunosuppressive including for after solid organ transplants, to prevent the activation of CD4- and CD8- T cells (Wilde & Goa 1996). However, the drug has since been discontinued due to the development of other treatments with similar efficacy and fewer side effects (Reichert 2012).</p>
Purity	>90 % by SDS PAGE
Flow Cytometry	Use 5 ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul
Storage	<p>Store at +4°C. DO NOT FREEZE.</p> <p>This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
Guarantee	Guaranteed until date of expiry. Please see product label.
Health And Safety Information	<p>Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA6146F</p> <p>10040</p>

Related Products

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL:FITC \(MCA929F\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M395440:220429'

Printed on 29 Apr 2024