

Datasheet: MCA6146GA

Description:	MOUSE ANTI HUMAN CD3
Specificity:	CD3
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	OKT3
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen	▪			
Functional Assays (1)	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) This product contains sodium azide, removal by dialysis is recommended prior to use in functional assays.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 0.5 mg/ml

Immunogen Human peripheral blood lymphocytes

External Database

Links

UniProt:

[P07766](#) [Related reagents](#)

Entrez Gene:

[916](#) CD3E [Related reagents](#)

Synonyms

T3E

Specificity

Mouse anti Human CD3 antibody, clone OKT3, recognizes T-cell surface glycoprotein CD3 epsilon chain (CD3 ϵ), also known as T-cell surface antigen T3/Leu-4 epsilon. CD3 ϵ is a subunit within the multiprotein cell surface receptor TCR-CD3 complex, which is expressed on the surface of mature T cells, NKT cells and thymocytes ([Mariuzza et al. 2020](#)). The CD3 complex is required for intracellular signaling and for TCR surface expression. Recruitment of the TCR/CD3 complex initiates early events leading to proliferation and differentiation of mature T cells into an effector cell ([Guy & Vignali 2009](#)).

Mice that lack the CD3 ϵ gene exhibit early arrest in T cell development and mutation of the gene in humans has been seen to result in severe combined immunodeficiency (SCID) ([DeJarnette et al. 1998](#), [de Saint Basile et al. 2004](#)).

Mouse anti Human CD3, clone OKT3 is commonly used in flow cytometry in the phenotyping human T cells. Characterization of the OKT3 clone has shown that it is able to compete and block the binding of the CD3 clones SK7 and UCHT1 to human T cells ([Li et al. 2005](#)). The OKT3 clone has also been found to have immunosuppressive properties in vivo. As result of this, the clone was humanized to be used as a therapeutic drug called Muromonomab (Orthoclone). Muromonomab was used as an immunosuppressive including for after solid organ transplants, to prevent the activation of CD4- and CD8- T cells ([Wilde & Goa 1996](#)). However, the drug has since been discontinued due to the development of other treatments with similar efficacy and fewer side effects ([Reichert 2012](#)).

Purity

>90 % by SDS PAGE

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee

12 months from date of despatch.

Health And Safety Information

Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M395441:220429'

Printed on 31 Oct 2022