

## Datasheet: MCA6150F BATCH NUMBER 159405

Description:	MOUSE ANTI PIG CD11c:FITC
Specificity:	CD11c
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	3A8
lsotype:	lgG1
Quantity:	0.1 mg

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u> rad-antibodies.com/protocols.							
		Yes	No	Not Determined	Suggested Dilution			
	Flow Cytometry	-			Neat - 1/10			
	Where this product ha	is not been tes	ted for u	se in a particular tecl	nnique this does not			
	necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.							
Target Species	Pig							
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid							
Max Ex/Em	Fluorophore	Excitation Ma	ıx (nm)	Emission Max (nm)				
	FITC	490		525				
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant							
Buffer Solution	Phosphate buffered saline							
Preservative Stabilisers	0.09% Sodium Azide (NaN <sub>3</sub> ) 1% Bovine Serum Albumin							
Approx. Protein Concentrations	IgG concentration 0.1	mg/ml						

Immunogen	Pig alveolar macrophages					
External Database Links	UniProt: K9IVW2 Related reagents					
Specificity	Mouse anti Pig CD11c, clone 3A8 recognizes CD11c also known as integrin alpha-X.					
	Porcine CD11c is mainly expressed by blood and skin polymorphonuclear and mononuclear phagocytes as well as monocytes, granulocytes and alveolar and inflammatory macrophages. Porcine CD11c is not expressed by plasmacytoid dendritic cells (pDC).					
	Mouse anti Pig CD11c, clone 3A8 antibody can be used to distinguish resident from migrating dendritic cells (DC) in lymph nodes;migrated DC express lower levels of surface CD11c and have higher MHC class II expression than resident DC. Both porcine cDC1 and cDC2 subsets express CD11c, with cDC1 appearing to have slightly lower levels than cDC2. Clone 3A8 did not show species cross-reactivity with human CD11c-CD18 transfectants or sheep myeloid cells (Deloizy <i>et al.</i> 2016).					
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul					
References	1. Álvarez, B. <i>et al.</i> (2023) Porcine Macrophage Markers and Populations: An Update. Cells. 12 (16) :2103.					
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.					
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.					
Guarantee	12 months from date of despatch					
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA6150F 10041					

# **Related Products**

## **Recommended Negative Controls**

### MOUSE IgG1 NEGATIVE CONTROL:FITC (MCA928F)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	d.com	Email: antibody_sales_uk@bio-ra	id.com	Email: antibody_sales_de@bio-rad.com

#### Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint