

Datasheet: MCA6161GA

Description:	RAT ANTI HUMAN C3d			
Specificity:	C3d			
Format:	Purified			
Product Type:	Monoclonal Antibody			
Clone:	3			
Isotype:	lgG2a			
Quantity:	0.1 mg			

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-						
	rad-antibodies.com/protocols.						
		Yes	No	Not Determined	Suggested Dilution		
	ELISA	•					
	Immunoprecipitation	-					
	Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.						
Target Species	Human						
Product Form	Purified IgG - liquid						
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant						
Buffer Solution	Phosphate buffered saline						
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)						
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml						
External Database Links	UniProt: P01024 Related	reagents					

	Entrez Gene: <u>718</u> C3 <u>Related reagents</u>			
Synonyms	CPAMD1			
Specificity	Rat anti Human C3d antibody, clone 3 (also known as clone YB2/39-11-7), recognizes a linear antigen on C3d. This antibody also detects C3, C3b, iC3b and C3dg, it does not recognize C3c. To distinguish C3 from C3b, a Mouse anti Human C3a/C3a desArg antibody (MCA6163GA) can be used.			
	C3 is a 185 kDa glycoprotein consisting of an alpha and beta chain and is the key component of the complement activation pathways. There are 3 activation pathways within the complement system; the classical, lectin and alternative pathways. All three pathways converge on C3, resulting in its cleavage to anaphylatoxin C3a and opsonin C3b. Once formed, C3b targets microbial cell surface receptors, marking them for phagocytosis by macrophages. The activity of C3b is heavily regulated to prevent accidental host tissue damage. Of these regulators, circulating Factor I, in the presence of co-factors, cleaves C3b to generate inactivated C3b (iC3b). iC3b's inactive designation relates to its inability to assist in the formation of convertases. In the presence of membrane bound complement receptor 1, iC3b is cleaved further to C3dg, before further cleavage into C3d and C3c. C3d is able to interact with complement receptor 2, which is expressed on the surface of B cells. C3d is involved in pathogen opsonisation, using a thiolester bond to covalently coat pathogens (Ricklin <i>et al.</i> 2016).			
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.			
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.			
Guarantee	12 months from date of despatch			
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA6161GA 10040			
Regulatory	For research purposes only			

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16)	DyLight®800
Rabbit Anti Rat IgG (STAR17)	<u>FITC</u>
Goat Anti Rat IgG (STAR73)	<u>RPE</u>
Rabbit Anti Rat IgG (STAR21)	<u>HRP</u>

Goat Ant	i Rat IgG (MOUSE ADSC	ORBED) (ST	[AR71) <u>DyLight®550</u> , [DyLight®650	DyLight®800
Goat Ant	i Rat IgG (STAR131)		Alk. Phos., Bio	<u>tin</u>	
Goat Ant	i Rat IgG (STAR72)		<u>HRP</u>		
Goat Ant	i Rat IgG (STAR69)		<u>FITC</u>		
North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad	d.com	Email: antibody_sales_uk@bio-	rad.com	Email: antibody_sales_de@bio-rad.com
To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M441983:250528'					
Printed on 28 May 2025					

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint