Datasheet: MCA6167GA BATCH NUMBER 163327

| Description: | MOUSE ANTI PIG CD33 |
|---------------|---------------------|
| Specificity: | CD33 |
| Format: | Purified |
| Product Type: | Monoclonal Antibody |
| Clone: | 5D5 |
| lsotype: | lgG1 |
| Quantity: | 0.1 mg |
| | |

Product Details

| Applications | This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> . | | | | | | | |
|-----------------------------------|--|-------------------|----|----------------|----------------------------|--|--|--|
| | | Yes | No | Not Determined | Suggested Dilution | | | |
| | Flow Cytometry | • | | | 1/50 - 1/100 | | | |
| | Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls. | | | | | | | |
| Target Species | Pig | | | | | | | |
| Product Form | Purified IgG - liquid | | | | | | | |
| Preparation | Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant | | | | | | | |
| Buffer Solution | Phosphate buffered salir | ne | | | | | | |
| Preservative Stabilisers | 0.09% Sodium Azide (Na | aN ₃) | | | | | | |
| Approx. Protein Concentrations | IgG concentration 1.0 m | g/ml | | | | | | |
| Immunogen | Recombinant protein co fused to the Fc portion o | - | | - | omains of porcine Siglec-3 | | | |

| Specificity | Mouse anti Pig CD33, clone 5D5 recognizes CD33, also known as Siglec-3. | | | | | |
|----------------------------------|--|--|--|--|--|--|
| | Sialic-acid-binding immunoglobulin-like lectin (Siglecs) are cell surface receptors belonging to the immunoglobulin superfamily and recognize terminal sialic acids present in complex oligosaccharides of glycoproteins or glycolipids. Siglecs are mainly expressed on the cells of the immune systems and play a regulatory role, modulating inflammatory and immune responses. | | | | | |
| | Clone 5D5 was used to detect Siglec-3 expression on monocytes (CD172a ⁺ or CD16 ^{hi} PBMCs), granulocytes (CD52 ⁺ and CD52 ⁻), and bone marrow. CD33 was also detected in lymph node, splenic, and alveolar macrophages, although at lower levels than on monocytes (<u>Alvarez <i>et al.</i> 2015).</u> | | | | | |
| Flow Cytometry | Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul | | | | | |
| References | Bohorquez, J.A. <i>et al.</i> (2019) Identification of an Immunosuppressive Cell Population during Classical Swine Fever Virus Infection and Its Role in Viral Persistence in the Host. <u>Viruses. 11 (9): 822.</u> Álvarez-Estrada, Á. <i>et al.</i> (2019) TLR2, Siglec-3 and CD163 expressions on porcine peripheral blood monocytes are increased during sepsis caused by <i>Haemophilus</i> <i>parasuis</i>. <u>Comp Immunol Microbiol Infect Dis. 64: 31-39.</u> Poderoso, T. <i>et al.</i> (2020) Expression of Siglec-1, -3, -5 and -10 in porcine cDC1 and cDC2 subsets from blood, spleen and lymph nodes and functional capabilities of these cells. <u>Dev Comp Immunol. 109: 103692.</u> Álvarez, B. <i>et al.</i> (2023) Porcine Macrophage Markers and Populations: An Update. <u>Cells. 12 (16): 2103.</u> | | | | | |
| Storage | This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. | | | | | |
| Guarantee | 12 months from date of despatch | | | | | |
| Health And Safety Information | Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA6167GA 10040 | | | | | |
| Regulatory | For research purposes only | | | | | |
| | | | | | | |

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)RPEGoat Anti Mouse IgG IgA IgM (STAR87...)HRP

| Goat Ant | i Mouse IgG (STAR76) | RPE | | | | |
|---|---|----------------|--|-----------------|--|--|
| Rabbit A | nti Mouse IgG (STAR13) | HR | - | | | |
| Goat Ant | i Mouse IgG (STAR70) | <u>FIT</u> | <u>C</u> | | | |
| Goat Ant | i Mouse IgG (H/L) (STAR11 | 7) <u>Alk.</u> | Phos., DyLight®488, Dy | Light®550, | | |
| | | DyL | ight®650, DyLight®680, | DyLight®80 | <u>D</u> , | |
| | | FIT | <u>C, HRP</u> | | | |
| Rabbit A | nti Mouse IgG (STAR9) | <u>FIT(</u> | <u>C</u> | | | |
| Goat Ant | i Mouse IgG (STAR77) | HR | | | | |
| Goat Anti Mouse IgG (Fc) (STAR120) <u>FITC, H</u> | | | <u>C, HRP</u> | | | |
| Recommended Negative Controls | | | | | | |
| MOUSE IgG1 NEGATIVE CONTROL (MCA928) | | | | | | |
| North & South America | Tel: +1 800 265 7376 W Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.co | orldwide m | Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad | Europe d.com | Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com | |

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M421903:230808'

Printed on 01 Feb 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint