

Datasheet: MCA6169GA

## **BATCH NUMBER 163340**

Description:	MOUSE ANTI PIG SIGLEC-10
Specificity:	SIGLEC-10
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	2E9
Isotype:	lgG1
Quantity:	0.1 mg

# **Product Details**

### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			1/200 - 1/1000

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Pig	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue cultu supernatant	re
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	0.09% Sodium Azide (NaN <sub>3</sub> )	
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml	
Immunogen	NIH/3T3 mouse cells transfected with plasmid pSiglec-10-GFP, encoding the posiglec-10 tagged with the green fluorescent protein (GFP) on the carboxy-terminates.	

#### **Specificity**

Mouse anti Pig Siglec-10, clone 2E9 recognizes Siglec-10.

Sialic-acid-binding immunoglobulin-like lectin (Siglecs) are cell surface receptors belonging to the immunoglobulin superfamily and recognize terminal sialic acids present in complex oligosaccharides of glycoproteins or glycolipids. Siglecs are mainly expressed on the cells of the immune systems and play a regulatory role, modulating inflammatory and immune responses. When expressed on transfected cells, porcine Siglec-10 is able to bind red blood cells in a sialic acid-dependent manner.

Mouse anti Pig Siglec-10 antibody, clone 2E9 has been successfully used to examine Siglec-10 cell and tissue distribution in swine. Siglec-10 was found to be expressed on blood B cells (CD21<sup>+</sup> or CD79a<sup>+</sup>) and in B cell areas of the spleen and lymph nodes. Weak expression was also detected on monocytes (<u>Escalona et al. 2014</u>).

## Flow Cytometry

Use 10ul of the suggested working dilution to label 1x10<sup>6</sup> cells in 100ul

#### References

- 1. Álvarez-Estrada, Á. *et al.* (2019) TLR2, Siglec-3 and CD163 expressions on porcine peripheral blood monocytes are increased during sepsis caused by *Haemophilus parasuis*. Comp Immunol Microbiol Infect Dis. 64: 31-39.
- 2. Poderoso, T. *et al.* (2019) Analysis of the expression of porcine CD200R1 and CD200R1L by using newly developed monoclonal antibodies. <u>Dev Comp Immunol. 100:</u> 103417.
- 3. Poderoso, T. *et al.* (2020) Expression of Siglec-1, -3, -5 and -10 in porcine cDC1 and cDC2 subsets from blood, spleen and lymph nodes and functional capabilities of these cells. <u>Dev Comp Immunol</u>. 109: 103692.
- 4. Álvarez, B. *et al.* (2023) Porcine Macrophage Markers and Populations: An Update. Cells. 12 (16): 2103.

### Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA6169GA">https://www.bio-rad-antibodies.com/SDS/MCA6169GA</a> 10040
Regulatory	For research purposes only

## **Related Products**

### **Recommended Secondary Antibodies**

Rabbit Anti Mouse IgG (STAR12...) RPE
Goat Anti Mouse IgG IgA IgM (STAR87...) HRP

Goat Anti Mouse IgG (STAR76...)

RPE
Goat Anti Mouse IgG (STAR70...)

FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Goat Anti Mouse IgG (STAR77...) HRP

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

Rabbit Anti Mouse IgG (STAR13...) HRP
Rabbit Anti Mouse IgG (STAR9...) FITC

# **Recommended Negative Controls**

### MOUSE IgG1 NEGATIVE CONTROL (MCA928)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M417822:230417'

### Printed on 23 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint