

Datasheet: MCA6181B

Description:	MOUSE ANTI HUMAN CCL17:Biotin
Specificity:	CCL17
Other names:	TARC
Format:	Biotin
Product Type:	Monoclonal Antibody
Clone:	I11-2A1
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunoassay	▪			1.0 ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG conjugated to Biotin - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	<i>E. coli</i> derived recombinant human full length CCL17 (Ala24-Ser94)

**External Database
Links**

UniProt:

[Q92583](#) [Related reagents](#)

Entrez Gene:

[6361](#) CCL17 [Related reagents](#)

Synonyms

SCYA17, TARC

Fusion Partners

Cell fusion between immunized BALB/c mouse spleen cells and mouse myeloma SP2/0

Specificity

Mouse anti Human CCL17 antibody, clone I11-2A1 recognizes C-C motif chemokine 17, also known as TARC and SCYA17. CCL17 is a C-C chemokine highly expressed by the thymus and other cell types, including fibroblasts ([Li et al. 2019](#)). CCL17 induces and enhances a broad spectrum of immune reactions, including some contact hypersensitivity reactions, allograft rejections, inflammatory bowel disease, atopic dermatitis and atherosclerosis ([Scheu et al. 2017](#)). Therapeutic use of antibodies to neutralize CCL17 can block influx of T cells and ameliorate the impact of various diseases ([Teplyakov et al. 2017](#)). Upregulation of CCL17 has been reported in breast cancer, and predicts a decrease in overall survival for patients with breast cancer ([Li et al. 2019](#)).

The biotinylated Mouse anti Human CCL17 antibody, clone I11-2A1 (MCA6181B) can be used as a detection antibody in a sandwich ELISA with the purified Mouse anti Human CCL17 antibody, clone I02-2A4 ([MCA6180GA](#)) as the capture antibody.

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee

12 months from date of despatch

**Health And Safety
Information**

Material Safety Datasheet documentation #10041 available at:
<https://www.bio-rad-antibodies.com/SDS/MCA6181B>
10041

Regulatory

For research purposes only

Related Products

Recommended Useful Reagents

[MOUSE ANTI HUMAN CCL17 \(MCA6180GA\)](#)

[RECOMBINANT HUMAN CCL17 \(PHP295\)](#)

**North & South
America**

Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M384485:210513'

Printed on 29 Aug 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)