

Datasheet: MCA618R

BATCH NUMBER 1807

Description:	MOUSE ANTI RAT CD11b
Specificity:	CD11b
Other names:	INTEGRIN ALPHA M CHAIN, MAC-1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	ED7
Isotype:	IgG1
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen (1)	▪			1/500
Immunohistology - Paraffin		▪		
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	
Functional Assays (2)	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

(2)This product contains sodium azide, removal by dialysis is recommended prior to use in functional assays. Bio-Rad recommend the use of [EQU003](#) for this purpose.

Target Species	Rat
Product Form	Purified IgG - liquid

Preparation	Purified IgG prepared by affinity chromatography on Protein A
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 0.5 mg/ml
Immunogen	Rat spleen cell homogenate.
RRID	AB_322901
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse Sp2/0 Ag-14 myeloma cell line.
Specificity	Mouse anti Rat CD11b antibody, clone ED7 recognizes a membrane antigen on rat macrophages, monocytes, dendritic cells and granulocytes. Like ED8, ED7 recognises small ramified microglia in the central nervous system. No other cell types are positive for ED7, except for the cilia of the bronchus epithelium. The recognised antigen is a heterodimer (~160 and ~95 kDa) belonging to the family of adhesion molecules (CD11b/CD18). ED7 and ED8 may recognise closely related epitopes on the same molecule. ED7 and ED8 have been shown to induce homotypic aggregation of granulocytes.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> 1. Damoiseaux, J.G. <i>et al.</i> (1989) Rat bone marrow and monocyte cultures: influence of culture time and lymphokines on the expression of macrophage differentiation antigens. J Leukoc Biol. 46 (3): 246-53. 2. Damoiseaux, J.G. <i>et al.</i> (1989) Heterogeneity of macrophages in the rat evidenced by variability in determinants: two new anti-rat macrophage antibodies against a heterodimer of 160 and 95 kd (CD11/CD18). J Leukoc Biol. 46 (6): 556-64. 3. DeGroot, C.J. <i>et al.</i> (1988) Discrimination between different types of neuroglial cells in rat central nervous system using combined immuno- and enzyme-histochemical methods. Immunobiology. 178 (3): 177-90. 4. Huitinga, I. <i>et al.</i> (1993) Treatment with anti-CR3 antibodies ED7 and ED8 suppresses experimental allergic encephalomyelitis in Lewis rats. Eur J Immunol. 23 (3): 709-15. 5. Drasković-pavlović, B. <i>et al.</i> (1999) Differential effects of anti-rat CD11b monoclonal antibodies on granulocyte adhesiveness. Immunology. 96 (1): 83-9. 6. Garn, H. <i>et al.</i> (2006) Phenotypical and functional characterization of alveolar macrophage subpopulations in the lungs of NO2-exposed rats. Respir Res. 7: 4. 7. Dick, A.D. <i>et al.</i> (2001) Distribution of OX2 antigen and OX2 receptor within retina. Invest Ophthalmol Vis Sci. 42 (1): 170-6. 8. Ramaglia, V. <i>et al.</i> (2007) The membrane attack complex of the complement system is essential for rapid Wallerian degeneration. J Neurosci. 27 (29): 7663-72.

Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA618R 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA1209\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M368720:200529'

Printed on 18 Jan 2024