

Datasheet: MCA620PE

Description:	MOUSE ANTI RAT CD172a:RPE
Specificity:	CD172a
Other names:	SIRP ALPHA
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	ED9
Isotype:	lgG1
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat - 1/5

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rat			
Product Form	Purified IgG conjug	ated to R. Phycoerythrin	(RPE) - lyophilized	
Reconstitution	Reconstitute with 1	ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)	
	RPE 488nm laser	496	578	
Preparation	Purified IgG prepar supernatant	ed by affinity chromatog	raphy on Protein A from tissue o	
Buffer Solution	Phosphate buffered	d saline		
Preservative	0.09% Sodium Azio	de		
Stabilisers	1% Bovine Serum Albumin			
	5% Sucrose			

Immunogen	Spleen cell homogenate.
External Database Links	UniProt: P97710 Related reagents
	Entrez Gene: 25528 Sirpa Related reagents
Synonyms	Bit, Mfr, Ptpns1, Shps1, Sirp
RRID	AB_322316
Fusion Partners	Spleen cells from immunised mice were fused with cells of the Sp2/0 Ag-14 myeloma cell line.
Specificity	Mouse anti Rat CD172a antibody, clone ED9 recognizes rat Tyrosine-protein phosphatase non-receptor type substrate 1, also known as CD172a, Signal-regulatory protein alpha-1, SIRPα -1, SHP substrate 1, Macrophage membrane protein MFP150 or Macrophage fusion receptor. CD172a is a 509 amino acid ~56 kDa single pass type 1 transmembrane glycoprotein expressed selectively by myeloid cells and by neurons (UniProt: P97710). Mouse anti Rat CD172a antibody, clone ED9 has been reported to bind to an alternative epitpe to another anti CD172 antibody, clone OX-41 (Adams et al. 1998) and has been reported to block the interaction of CD172a - CD47 (de Vries et al. 2002).
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	1. Damoiseaux, J.G. <i>et al.</i> (1989) Rat bone marrow and monocyte cultures: influence of culture time and lymphokines on the expression of macrophage differentiation antigens. <u>J Leukoc Biol. 46 (3): 246-53.</u>
	2. Damoiseaux, J.G. <i>et al.</i> (1989) Heterogeneity of macrophages in the rat evidenced by variability in determinants: two new anti-rat macrophage antibodies against a heterodimer of 160 and 95 kd (CD11/CD18). <u>J Leukoc Biol. 46 (6): 556-64.</u>
	3. Adams, S. <i>et al.</i> (1998) Signal-regulatory protein is selectively expressed by myeloid and neuronal cells. <u>J Immunol. 161 (4): 1853-9.</u>
	4. DeVries, H.E. <i>et al.</i> (2002) Signal-regulatory protein alpha-CD47 interactions are required for the transmigration of monocytes across cerebral endothelium. <u>J Immunol. 168</u> (11): 5832-9.
	5. Blackbeard, J. et al. (2007) Quantification of the rat spinal microglial response to
	navinhaval namia injumi na vasta alad hujimansunahiata ahamiaal imaana analysia anad flass

- 5. Blackbeard, J. *et al.* (2007) Quantification of the rat spinal microglial response to peripheral nerve injury as revealed by immunohistochemical image analysis and flow cytometry. <u>J Neurosci Methods</u>. 164 (2): 207-17.
- 6. Bode, U. *et al.* (2008) Dendritic cell subsets in lymph nodes are characterized by the specific draining area and influence the phenotype and fate of primed T cells. Immunology. 123 (4): 480-90.
- 7. Chang, J.C. *et al.* (2019) Early Immune Response to Acute Gastric Fluid Aspiration in a Rat Model of Lung Transplantation. <u>Exp Clin Transplant</u>. 17 (1): 84-92.
- 8. Holler, J. et al. (2008) Neuropeptide Y is expressed by rat mononuclear blood

leukocytes and strongly down-regulated during inflammation. <u>J Immunol. 181 (10):</u> 6906-12.

9. Schmiedl, A. *et al.* (2021) Lung development and immune status under chronic LPS exposure in rat pups with and without CD26/DPP4 deficiency. <u>Cell Tissue Res. Oct 04 [Epub ahead of print].</u>

Storage

Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA620PE 20487
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL: RPE (MCA1209PE)

North & South Tel: +1 800 265 7376

America

Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M419501:230616'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint