

Datasheet: MCA6308

BATCH NUMBER 168754

Description:	RABBIT ANTI ANDROGEN RECEPTOR
Specificity:	ANDROGEN RECEPTOR
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	RM254
Isotype:	IgG
Quantity:	0.1 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Paraffin	▪			1/1000 - 1/2500
Western Blotting	▪			1/100 - 1/1000

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography of Protein A from animal origin-free culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide
Stabilisers	1% Bovine Serum Albumin 50% Glycerol
Immunogen	A peptide corresponding to the N-terminus of androgen receptor
External Database	UniProt:

Links

[P10275](#) [Related reagents](#)

Entrez Gene:

[367](#) AR [Related reagents](#)

Synonyms

DHTR, NR3C4

Specificity

Rabbit anti Human androgen receptor antibody recognizes androgen receptor (AR), also known as Dihydrotestosterone receptor (DHTR), NR3C4. Androgen Receptor is a ligand-activated transcription factor that belongs to a large family of nuclear receptor proteins. Unbound Androgen Receptor resides predominately in the cytoplasm as a heteromeric complex with hsp90 and other chaperone proteins. Circulating testosterone is converted to dihydrotestosterone (DHT) which binds to androgen receptor in the cytoplasm, releasing androgen receptor from its inhibition with HSPs and allowing translocation to the nucleus, where androgen receptor can initiate transcription of its target genes ([Tan et al. 2015](#)). Androgen signaling plays a role in pathogenesis of prostate cancer, and gonadal depletion of androgens is a central therapy for advanced prostate cancer. However, progression to castration-resistant prostate cancer has led to development of therapies that directly target the androgen receptor ([Dai et al. 2017](#)) Androgen receptor also seems to play a major role in carcinogenesis of triple negative breast cancer (TNBC), a very aggressive cancer type which comprises 15 to 20% of all breast cancers ([Gerratana et al. 2018](#)).

Storage

Store at -20°C only.
Storage in frost-free freezers is not recommended.
This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10048 available at: <https://www.bio-rad-antibodies.com/SDS/MCA6308>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)

Sheep Anti Rabbit IgG (STAR35...) [RPE](#)

Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M376287:210127'

Printed on 13 Oct 2025

