

# Datasheet: MCA6308

Description:	RABBIT ANTI ANDROGEN RECEPTOR
Specificity:	ANDROGEN RECEPTOR
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	RM254
lsotype:	lgG
Quantity:	0.1 ml

#### **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
		Yes	No	Not Determined	Suggested Dilution	
	Immunohistology - Paraffin	-			1/1000 - 1/2500	
	Western Blotting	•			1/100 - 1/1000	
	Where this product has r	not been te	ested for	use in a particular tech	nique this does not	
	necessarily exclude its us a guide only. It is recomn system using appropriate	nended th	at the use	er titrates the product f		
Target Species	Human					
Product Form	Purified IgG - liquid					
Preparation	Purified IgG prepared by culture supernatant	affinity ch	nromatogr	raphy of Protein A from	n animal origin-free	
Buffer Solution	Phosphate buffered salin	е				
Preservative	0.09% Sodium Azide					
Stabilisers	1% Bovine Serum Album	nin				
	50% Glycerol					
Immunogen	A peptide corresponding	to the N-t	erminus o	of androgen receptor		
External Database Links	UniProt:					

	P10275 Related reagents
	Entrez Gene:
	<u>367</u> AR <u>Related reagents</u>
Synonyms	DHTR, NR3C4
Specificity	<b>Rabbit anti Human androgen receptor antibody</b> recognizes androgen receptor (AR), also known as Dihydrotestosterone receptor (DHTR), NR3C4. Androgen Receptor is a ligand-activated transcription factor that belongs to a large family of nuclear receptor proteins. Unbound Androgen Receptor resides predominately in the cytoplasm as a heteromeric complex with hsp90 and other chaperone proteins. Circulating testosterone is converted to dihydrotestosterone (DHT) which binds to androgen receptor in the cytoplasm, releasing androgen receptor from its inhibition with HSPs and allowing translocation to the nucleus, where androgen receptor can initiate transcription of its target genes (Tan et al. 2015). Androgen signaling plays a role in pathogenesis of prostate cancer, and gonadal depletion of androgens is a central therapy for advanced prostate cancer. However, progression to castration-resistant prostate cancer has led to development of therapies that directly target the androgen receptor (Dai et al. 2017) Androgen receptor also seems to play a major role in carcinogenesis of triple negative breast cancer (TNBC), a very aggressive cancer type which comprises 15 to 20% of all breast cancers (Gerratana et al. 2018).
Storage	Store at -20°C only. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10048 available at: https://www.bio-rad-antibodies.com/SDS/MCA6308 10048
Regulatory	For research purposes only

## **Related Products**

### **Recommended Secondary Antibodies**

Sheep Anti Rabbit IgG (STAR34)	<u>FITC</u>
Goat Anti Rabbit IgG (H/L) (STAR124	) <u>HRP</u>
Goat Anti Rabbit IgG (Fc) (STAR121)	Biotin, FITC, HRP
Sheep Anti Rabbit IgG (STAR35)	RPE
Sheep Anti Rabbit IgG (STAR36)	DyLight®488, DyLight®680, DyLight®800

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad	l.com	Email: antibody_sales_uk@bio-ra	ad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M376287:210127'

#### Printed on 29 Aug 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint