

Datasheet: MCA6368

Description:	RABBIT ANTI CD278
Specificity:	CD278
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	RM417
Isotype:	IgG
Quantity:	0.1 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Paraffin	▪			1/100 - 1/200

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from animal origin-free culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide
Stabilisers	1% Bovine Serum Albumin 50% Glycerol
Immunogen	A peptide corresponding to the C-terminus of human CD278

External Database

Links

UniProt:

[Q9Y6W8](https://www.uniprot.org/entry/Q9Y6W8)

[Related reagents](#)

Entrez Gene:[29851](#) ICOS [Related reagents](#)

Synonyms AILIM

Specificity **Rabbit anti CD278 antibody** recognizes inducible T-cell costimulator, also known as ICOS and AILIM.

CD278 is a member of the immunoglobulin family of co-receptor molecules mainly expressed on activated cytotoxic T cells, memory T cells, and regulatory T cells ([Xiao et al. 2020](#)). Its main role appears to be supporting differentiation and function of CD4+ follicular helper (Tfh) cells during germinal centre reactions ([Panneton et al. 2018](#)). The protein has significant homology to CD28 and CTLA-4. CD278 expression is induced on T cells following T cell receptor (TCR) cross-linking and/or CD28 co-stimulation ([Wikenheiser and Stumhofer 2016](#)). Because of its expression pattern, CD278 provides an early and sensitive method for detecting T-cell immune responses to cancer therapy ([Xiao et al. 2020](#)).

Storage Store at -20°C only.
Storage in frost-free freezers is not recommended.
This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10048 available at: <https://www.bio-rad-antibodies.com/SDS/MCA6368>
10048

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...) [FITC](#)Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)Sheep Anti Rabbit IgG (STAR35...) [RPE](#)Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)Sheep Anti Rabbit IgG (STAR36...) [DyLight®488](#), [DyLight®680](#), [DyLight®800](#)**North & South** Tel: +1 800 265 7376**America** Fax: +1 919 878 3751Email: antibody_sales_us@bio-rad.com**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M387000:210604'

Printed on 25 Mar 2023

