

Datasheet: MCA637GA

Description:	MOUSE ANTI PIG IgM
Specificity:	IgM
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	K52 1C3
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA	▪			1/5000 - 1/100,000
Immunoprecipitation			▪	
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Pig
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Porcine IgM
Fusion Partners	Spleen cells of immunised mice were fused with cells of the P3 - X63 - Ag 8.653 mouse myeloma

line.

Specificity	Mouse anti Pig IgM antibody, clone K52 1C3 recognizes porcine IgM heavy chain. No cross-reactivity with porcine IgA and IgG is seen in ELISA.
--------------------	--

Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul
-----------------------	--

References	<ol style="list-style-type: none">1. Leitão, A. <i>et al.</i> (2001) The non-haemadsorbing African swine fever virus isolate ASFV/NH/P68 provides a model for defining the protective anti-virus immune response. J Gen Virol. 82 (Pt 3): 513-23.2. Baltes, N. <i>et al.</i> (2001) Actinobacillus pleuropneumoniae iron transport and urease activity: effects on bacterial virulence and host immune response. Infect Immun. 69 (1): 472-8.3. Hamano, M. <i>et al.</i> (2007) Detection of antibodies to Japanese encephalitis virus in the wild boars in Hiroshima prefecture, Japan. Epidemiol Infect. 135: 974-7.4. Andersen, J.K. <i>et al.</i> (1999) Systematic characterization of porcine ileal Peyer's patch, I. apoptosis-sensitive immature B cells are the predominant cell type. Immunology. 98 (4): 612-21.5. Bailey, M. (2004) Effects of infection with transmissible gastroenteritis virus on concomitant immune responses to dietary and injected antigens. Clin Diagn Lab Immunol. 11: 337-43.6. Pasternak, J.A. <i>et al.</i> (2015) Oral antigen exposure in newborn piglets circumvents induction of oral tolerance in response to intraperitoneal vaccination in later life. BMC Vet Res. 11: 350.7. Seele, J. <i>et al.</i> (2015) The immunoglobulin M-degrading enzyme of Streptococcus suis, IdeSsuis, is a highly protective antigen against serotype 2. Vaccine. 33 (19): 2207-12.8. Stepanova, H. <i>et al.</i> (2011) Association of attenuated mutants of <i>Salmonella enterica</i> serovar Enteritidis with porcine peripheral blood leukocytes. FEMS Microbiol Lett. 321: 37-42.9. Laycock, G. <i>et al.</i> (2012) A defined intestinal colonization microbiota for gnotobiotic pigs. Vet Immunol Immunopathol. 149: 216-24.10. Lewis MC <i>et al.</i> (2013) Dietary supplementation with Bifidobacterium lactis NCC2818 from weaning reduces local immunoglobulin production in lymphoid-associated tissues but increases systemic antibodies in healthy neonates. Br J Nutr. 110: 1243-52.11. Chen, F. <i>et al.</i> (2015) Generation of B Cell-Deficient Pigs by Highly Efficient CRISPR/Cas9-Mediated Gene Targeting. J Genet Genomics. 42 (8): 437-44.12. Rahe, M.C. & Murtaugh, M.P. (2017) Interleukin-21 Drives Proliferation and Differentiation of Porcine Memory B Cells into Antibody Secreting Cells. PLoS One. 12 (1): e0171171.13. Buermann, A. <i>et al.</i> (2018) Pigs expressing the human inhibitory ligand PD-L1 (CD 274) provide a new source of xenogeneic cells and tissues with low immunogenic properties. Xenotransplantation. 25 (5): e12387.
-------------------	---

Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody.
----------------	--

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
 Goat Anti Mouse IgG (STAR77...) [HRP](#)
 Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
 Rabbit Anti Mouse IgG (STAR8...) [DyLight@800](#)
 Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
 Goat Anti Mouse IgG (STAR76...) [RPE](#)
 Goat Anti Mouse IgG (STAR70...) [FITC](#)
 Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
 Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
 Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight@488](#), [DyLight@680](#),
[DyLight@800](#), [FITC](#), [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M368744:200529'

Printed on 11 Aug 2020

© 2020 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)