

Datasheet: MCA6398GA

### **BATCH NUMBER 162382**

Description:	MOUSE ANTI HORSE CD25
Specificity:	CD25
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	15-1
Isotype:	lgG1
Quantity:	0.1 mg

## **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			1/25 - 1/50

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Horse	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein A supernatant	A from tissue culture
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	0.09% Sodium Azide (NaN <sub>3</sub> )	
Carrier Free	Yes	
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml	
Immunogen	Recombinant equine CD25	

External Database Links	UniProt:  A8VUE0 Related reagents			
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the X3.Ag8-653 myeloma cell line			
Specificity	Mouse anti Horse CD25, clone 15-1 recognizes horse CD25, also known as the interleukin-2 (IL-2) receptor subunit alpha.			
	CD25 is expressed by activated T and B cells ( <u>Jackson et al. 1990</u> , <u>Butcher et al. 1990</u> ). CD25 expression on CD4+ T cells is used to identify regulatory T (Treg) cells. CD25 acts as the low affinity receptor for IL-2, but it also assembles into a heterotrimer with CD122 and CD132 to form the high affinity receptor.			
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul			
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.			
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.			
Guarantee	12 months from date of despatch			
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA6398GA">https://www.bio-rad-antibodies.com/SDS/MCA6398GA</a> 10040			
Regulatory	For research purposes only			

# **Related Products**

## **Recommended Secondary Antibodies**

Goat Anti Mouse IgG (STAR77...)

Rabbit Anti Mouse IgG (STAR12...)

RPE

Goat Anti Mouse IgG (STAR70...)

FITC

Goat Anti Mouse IgG IgA IgM (STAR87...) Alk. Phos., HRP

Goat Anti Mouse IgG (STAR76...) RPE

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Rabbit Anti Mouse IgG (STAR13...) HRP

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

Rabbit Anti Mouse IgG (STAR9...) FITC

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

Fax: +1 919 878 3751 Fax: +44 (0)1865 852 739 Fax: +49 (0) 89 8090 95 50 To Email: antibody\_sales\_us@bio-rad.com Email: antibody\_sales\_uk@bio-rad.com Email: antibody\_sales\_de@bio-rad.com Email: antibody\_sales\_de@bio-rad.com

batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M403816:230124'

#### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint