

## Datasheet: MCA6434GA

Description:	MOUSE ANTI HUMAN CYTOKERATIN (PAN REACTIVE)
Specificity:	CYTOKERATIN PAN REACTIVE
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	C-11
Isotype:	lgG1
Quantity:	0.1 mg

# **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further						
	rad-antibodies.com/proto	information. For general protocol recommendations, please visit <u>www.bio-</u>					
		Yes	No	Not Determined	Suggested Dilution		
	Flow Cytometry (1)	•			1/100		
	Immunohistology - Frozen						
	Immunohistology - Paraffin (2)	-					
	Immunoprecipitation						
	Western Blotting	•					
	Immunofluorescence						
	Rare Cell / CTC Enumeration	-			1/200		
	<ul> <li>Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.</li> <li>(1)Membrane permeabilization is required for this application. The use of Leucoperm (Product Code <u>BUF09</u>) is recommended for this purpose.</li> <li>(2)This product requires protein digestion pre-treatment of paraffin sections e.g. trypsin or pronase.</li> </ul>						
Target Species	Human						
Species Cross Reactivity	Based on sequence similarity, is expected to react with:Mammals <b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.						

Product Form	Purified IgG - liquid					
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant					
Buffer Solution	Phosphate buffered saline					
Preservative Stabilisers	<0.1% Sodium Azide (NaN <sub>3</sub> )					
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml					
Immunogen	Keratin-enriched preparation from human epidermoid carcinoma cell line A431					
External Database Links	UniProt: P02538 Related reagents					
	P05783 Related reagents					
	P13645 Related reagents					
	P13647 Related reagents					
	P13646 Related reagents					
	P19013 Related reagents					
	P48668 Related reagents					
	P04259 Related reagents					
	P05787 Related reagents					
	Entrez Gene:					
	<u>3853</u> KRT6A <u>Related reagents</u>					
	3875 KRT18 Related reagents					
	<u>3851</u> KRT4 <u>Related reagents</u>					
	3852 KRT5 Related reagents					
	3858 KRT10 Related reagents					
	3860 KRT13 Related reagents					
	<u>3854</u> KRT6B <u>Related reagents</u>					
	3856 KRT8 Related reagents					
	286887 KRT6C Related reagents					
Synonyms	CYK18, CYK4, CYK8, K6A, K6B, KPP, KRT6D, KRT6E, KRTL1					
Specificity	Mouse anti human cytokeratin (Pan reactive) antibody, clone C-11 recognizes the human cytokeratins 4, 5, 6, 8, 10, 13 and 18.					
	Cytokeratins, also known as keratins, form a subfamily of cytoplamsic proteins expressed in epithelial cells where they form intermediate filaments responsible for providing cells resistance to mechanical stress. Cytokeratins are characterized by their remarkable biochemical diversity, with epithelial tissues expressing at least 20 different cytokeratins					

	(cytokeratin 1 to 20) divided into type I cytokeratins (acidic cytokeratins 9 to 20) and type II cytokeratins (basic to neutral cytokeratins 1 to 8) subfamilies. Cytokeratin expression profile is largely specific to particular organs and tissues and their development stage. That tissue-specific expression profile of cytokeratins is usually retained upon epithelial cell transformation and is therefore used in clinic to identify the cells from which carcinomas originated from.
	Mouse anti Human Cytokeratin (Pan reactive) antibody, clone C-11, has been validated for use on the <u>Genesis Cell Isolation System with the CelSelect Slide<sup>TM</sup> technology</u> .
Flow Cytometry	Use 10ul of the suggested working dilution to label $1 \times 10^6$ cells in 100ul.
References	<ol> <li>Bártková, J.<i>et al.</i> (1991) Effects of tissue fixation conditions and protease pretreatment on immunohistochemical performance of a large series of new anti-keratin monoclonal antibodies: value in oncopathology. <u>Neoplasma. 38:439-446.</u></li> <li>Hoefler, R.S. &amp; , Kudva, I.T. (2021) EDL933 Strains of Escherichia coli O157 can Demonstrate Genetic Diversity and Differential Adherence to Bovine Recto-Anal Junction Squamous Epithelial Cells <u>The Open Microbiology Journal. 15 (1): 129-38.</u></li> <li>Kudva, I.T. <i>et al.</i> (2023) Bovine Rectoanal Junction <i>In Vitro</i> Organ Culture Model System to Study Shiga Toxin-Producing <i>Escherichia coli</i> Adherence. <u>Microorganisms. 11</u> (5): 1289.</li> </ol>
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch.
Acknowledgements	CelSelect Slides is a trademark of Bio-Rad Laboratories, Inc. in certain jurisdictions
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA6434GA 10040
Regulatory	For research purposes only

## **Related Products**

## **Recommended Secondary Antibodies**

Rabbit Anti Mouse IgG (STAR12)	RPE			
Goat Anti Mouse IgG IgA IgM (STAR87) <u>HRP</u>				
Goat Anti Mouse IgG (STAR76)	RPE			
Rabbit Anti Mouse IgG (STAR13)	HRP			
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>			
Goat Anti Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLight®550,			
	DyLight®650, DyLight®680, DyLight®800,			
	FITC, HRP			

Rabbit Anti Mouse IgG (STAR9...) FITC

Goat Anti Mouse IgG (STAR77...) HRP

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

### **Recommended Negative Controls**

### MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	ad.com	Email: antibody_sales_uk@bio-ra	ad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M426092:231120'

#### Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint