

Datasheet: MCA643PE

| | |
|----------------------|-------------------------|
| Description: | MOUSE ANTI RAT CD44:RPE |
| Specificity: | CD44 |
| Other names: | H-CAM, PGP-1 |
| Format: | RPE |
| Product Type: | Monoclonal Antibody |
| Clone: | OX-50 |
| Isotype: | IgG1 |
| Quantity: | 100 TESTS |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry | ▪ | | | Neat |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

| | | | |
|------------------------|---|----------------------------|--------------------------|
| Target Species | Rat | | |
| Product Form | Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized | | |
| Reconstitution | Reconstitute with 1 ml distilled water | | |
| Max Ex/Em | Fluorophore | Excitation Max (nm) | Emission Max (nm) |
| | RPE 488nm laser | 496 | 578 |
| Preparation | Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant | | |
| Buffer Solution | Phosphate buffered saline | | |
| Preservative | 0.09% Sodium Azide | | |
| Stabilisers | 1% | Bovine Serum Albumin | |
| | 5% | Sucrose | |

| | |
|--------------------------------|--|
| Immunogen | Rat T cell blasts. |
| External Database Links | <p>UniProt: P26051 Related reagents</p> <p>Entrez Gene: 25406 Cd44 Related reagents</p> |
| RRID | AB_321723 |
| Fusion Partners | Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS0/1 myeloma cell line. |
| Specificity | <p>Mouse anti Rat CD44 antibody, clone OX-50 recognizes the rat CD44 cell surface antigen, also known as Extracellular matrix receptor III, P90 lymphocyte homing/adhesion receptor, HUTCH-I, Hermes antigen, Hyaluronate receptor, Phagocytic glycoprotein 1, PGP-1 or Phagocytic glycoprotein I.</p> <p>CD44 is a 482 amino acid ~85 kDa single pass type I transmembrane glycoprotein, expressed by T cells, B cells, macrophages and thymocytes, with expression being increased following activation.</p> |
| Flow Cytometry | Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul. |
| References | <ol style="list-style-type: none"> 1. Stevenson, K.S. <i>et al.</i> (2009) Isolation, characterization, and differentiation of thy1.1-sorted pancreatic adult progenitor cell populations. Stem Cells Dev. 18:1389-98. 2. Jiang, T.S. <i>et al.</i> (2010) Reconstruction of the corneal epithelium with induced marrow mesenchymal stem cells in rats. Mol Vis. 16: 1304-16. 3. Kanellis, J. <i>et al.</i> (2010) JNK signalling in human and experimental renal ischaemia/reperfusion injury. Nephrol Dial Transplant. 25: 2898-908. 4. Li, S. <i>et al.</i> (2010) Upregulation of CXCR4 favoring neural-like cells migration via AKT activation. Neurosci Res. 67: 293-9. 5. Stephens, L.A. <i>et al.</i> (2004) Phenotypic characterization of regulatory CD4+CD25+ T cells in rats. Int Immunol. 16: 365-75. 6. Rice, C.M. <i>et al.</i> (2010) Multipotent adult progenitor cell isolation and proliferation in cytokine and serum-free medium conditioned by rat B104 cells. Br J Haematol. 148: 441-4. 7. Carulli, D. <i>et al.</i> (2006) Composition of perineuronal nets in the adult rat cerebellum and the cellular origin of their components. J Comp Neurol. 494: 559-77. 8. Galtrey, C.M. <i>et al.</i> (2008) Distribution and synthesis of extracellular matrix proteoglycans, hyaluronan, link proteins and tenascin-R in the rat spinal cord. Eur J Neurosci. 27: 1373-90. 9. Hofmann, N. <i>et al.</i> (2002) Increased expression of ICAM-1, VCAM-1, MCP-1, and MIP-1 alpha by spinal perivascular macrophages during experimental allergic encephalomyelitis in rats. BMC Immunol. 3: 11. 10. Walther, M. <i>et al.</i> (2001) Exogenous antigen containing perivascular phagocytes induce a non-encephalitogenic extravasation of primed lymphocytes. J Neuroimmunol. |

[117: 30-42.](#)

11. Suzuki, A. *et al.* (2006) Localization of CD44 and hyaluronan in the synovial membrane of the rat temporomandibular joint. [Anat Rec A Discov Mol Cell Evol Biol. 288: 646-52.](#)
12. Goransson, V. *et al.* (2004) Renal hyaluronan accumulation and hyaluronan synthase expression after ischaemia-reperfusion injury in the rat. [Nephrol Dial Transplant. 19: 823-30.](#)
13. Campbell, N.G. *et al.* (2016) Cell Size Critically Determines Initial Retention of Bone Marrow Mononuclear Cells in the Heart after Intracoronary Injection: Evidence from a Rat Model. [PLoS One. 11 \(7\): e0158232.](#)
14. Bejar, M.T. *et al.* (2016) Inhibition of Notch rescues the angiogenic potential impaired by cardiovascular risk factors in epicardial adipose stem cells. [FASEB J. 30 \(8\): 2849-59.](#)
15. Rochefort, G.Y. *et al.* (2006) Multipotential mesenchymal stem cells are mobilized into peripheral blood by hypoxia. [Stem Cells. 24 \(10\): 2202-8.](#)
16. Redondo, J. *et al.* (2015) Reductions in kinesin expression are associated with nitric oxide-induced axonal damage. [J Neurosci Res. 93 \(6\): 882-92.](#)
17. Huang, X. *et al.* (2019) MRI Tracking of SPIO- and *Fth1*-Labeled Bone Marrow Mesenchymal Stromal Cell Transplantation for Treatment of Stroke. [Contrast Media Mol Imaging. 2019: 5184105.](#)
18. Aminzadeh, A. *et al.* (2020) Investigating The Alterations of Oxidative Stress Status, Antioxidant Defense Mechanisms, MAP Kinase and Mitochondrial Apoptotic Pathway in Adipose-Derived Mesenchymal Stem Cells from STZ Diabetic Rats. [Cell J. 22 \(Suppl 1\): 38-48.](#)
19. Paiva, R.G. *et al.* (2020) Stem cells in end-to-side neurorrhaphy. Experimental study in rats [Acta Cirúrgica Brasileira. 35 \(12\) \[Epub ahead of print\].](#)
20. Hou, B. *et al.* (2018) Comparison of the Effects of BMSC-derived Schwann Cells and Autologous Schwann Cells on Remyelination Using a Rat Sciatic Nerve Defect Model. [Int J Biol Sci. 14 \(13\): 1910-22.](#)
21. Porwal, K. *et al.* (2019) Increased bone marrow-specific adipogenesis by clofazimine causes impaired fracture healing, osteopenia and osteonecrosis without extra-skeletal effects in rats. [Toxicol Sci. kfz172.](#)
22. Chang, H.H. *et al.* (2019) Intrarenal Transplantation of Hypoxic Preconditioned Mesenchymal Stem Cells Improves Glomerulonephritis through Anti-Oxidation, Anti-ER Stress, Anti-Inflammation, Anti-Apoptosis, and Anti-Autophagy. [Antioxidants \(Basel\). 9 \(1\): 2.](#)
23. Wu, J. *et al.* (2022) Reinforcing the function of bone graft via the Ca-P ceramics dynamic behavior-enhanced osteogenic microenvironment for optimal bone regeneration and reconstruction [Applied Materials Today. 27: 101465.](#)
24. Cheng, Y.H. *et al.* (2022) Intrarenal Arterial Transplantation of Dexmedetomidine Preconditioning Adipose Stem-Cell-Derived Microvesicles Confers Further Therapeutic Potential to Attenuate Renal Ischemia/Reperfusion Injury through miR-122-5p/Erythropoietin/Apoptosis Axis. [Antioxidants \(Basel\). 11\(9\): 1702.](#)

Storage

Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #20487 available at: <https://www.bio-rad-antibodies.com/SDS/MCA643PE>
20487

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:RPE \(MCA1209PE\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M419641:230616'

Printed on 14 Apr 2025

© 2025 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)